MODERN MEDICINE AND PRACTICE
http://inovatus.es/index.php/ejmmp

EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 4 No. 4 (Apr - 2024) EJMMP ISSN: 2795-921X

Physiological Changes Associated with Untreated Hypertension in Diwaniyah Province

Hanaa Abdul Abbas
Al-Qadisiya University, College of Pharmacy, Department of Clinical and Laboratory Science
hanaa.abdulameer@qu.edu.iq

Neeran Faleh Hassan
Al-Qadisiya University, College of Pharmacy, Department of Clinical and Laboratory Science

Ahmed Riyad Hamza
Al-Qadisiya University, College of Pharmacy, Department of Clinical and Laboratory Science

Abstract:
Hypertension is a disorder that is defined by raised blood pressure, and this study highlights the need of future research to better our knowledge of hypertension. As a result of the pumping activity of the heart, blood pressure is the force that is exerted by the blood that is circulating on the walls of the blood vessels. The cardiac output is multiplied by the systemic vascular resistance to arrive at the estimate. It is possible for people who have arterial hypertension to have enhanced cardiac output, elevated systemic vascular resistance, or both of these symptoms. Enhanced α-adrenoceptor activity or an increased release of peptides like angiotensin and endothelins may be ascribed to the rise in vascular tone. These peptides eventually increase the amount of cytosolic calcium in the smooth muscle of the vascular system, resulting in vasoconstriction. Furthermore, these peptides are responsible for the facilitation of vascular remodeling, which is a thickening of the smooth muscle of the vascular system. Left ventricular hypertrophy and diastolic dysfunction are two possible outcomes that might result from increased systemic vascular resistance and vascular stiffness. These factors exert greater load on the left ventricle. Several organs, including the heart, brain, kidneys, and eyes, are susceptible to experiencing harmful effects as a result of these alterations. If the measurement of your blood pressure is higher than 180/120 mm Hg, you are in a serious emergency situation that requires immediate medical attention. Because to white coat hypertension, a person's blood pressure ranges from low to high throughout the day and may even increase during medical examinations. As a result, obtaining an accurate diagnosis sometimes requires taking many readings over the course of several visits. Controlling high blood pressure requires a methodical and progressive approach to therapy, with the goal of bringing the diastolic blood pressure down to 100 mmHg over the course of 48–72 hours. It is essential to maintain control of hypertension by diet and continuing medication. Rapid declines in blood pressure may be harmful since brain arteries react slowly to changes. Sudden drops in blood pressure can also induce a stroke or other consequences.

Key words: Hypertension, Consequences, Heart Disease, Blood Pressure

Introduction

High blood pressure, often known as hypertension, is a potentially life-threatening health condition that develops when the force of the blood on the walls of the arteries is consistently too demanding. Hypertension is more frequently known as high blood pressure. The presence of high blood pressure is a defining feature of this condition. It is one of the characteristics that sets this illness apart from others, and that is the fact that the person has a high blood pressure. In the process of delivering oxygen and nutrients to the different organs in the body, blood pressure is an important component that must be carefully monitored and maintained. This is because blood pressure is a component that is crucial to the process. Because blood pressure is a component of the process that involves the circulation of blood via the circulatory system, this is the reason why this is the case. Because of this, this is the reason why it occurs. The reason for this is because arteries, which are the channels that transport blood out from the heart and to the organs of the body, are analogous to hoses. This is the reason why this is something that happens. The reason for the occurrence of this specific phenomena may be explained as follows. This is the reason why things transpire in the manner that they do. instances of the same thing are the wide artery and the small nozzle of a garden hose. Both of these are instances of circulation. A wide artery is akin to a garden hose, but a smaller artery is equivalent to the narrow nozzle of a garden hose. This is because of the similarities between the two. Due to the fact that the nozzle was altered to be more restrictive, there was an increase in the pressure that was present inside the hose. All of this occurred as a consequence of the nozzle being made more restrictive. According to the results of Egan et al. (2010), this phenomenon is analogous to the increase in arterial pressure that occurs inside the body when capillaries become constricted. This phenomenon is equivalent to the rise in arterial pressure. "mmHg" (millimeters of mercury) is the unit of measurement, thus the results of a blood pressure test are often stated in the form of two numbers, such as 120/70 mmHg. This is because "mmHg" is the unit of measurement. In addition to that, the information may be given in the form of a relation. Considering that the measurement of blood pressure is carried out in millimeters of mercury, this is the reason why this is the situation that has arisen. The pressure that is exerted by the heart as it contracts and pumps blood back and forth is referred to as the systolic blood pressure, and this is the pressure that is being measured here. The systolic arterial blood pressure constitutes the first number in the equation. The SYSTOLIC version of the blood pressure should be the first value that is shown. The measurements of the systolic blood pressure give evidence of this pressure; they show that this pressure is present in the body. While the heart is at rest between beats, the diastolic pressure, which is the second number in the equation, provides an indicator of the pressure that is present in the heart. It is the diastolic pressure that is responsible for this pressure. On this particular occurrence, the amount of pressure that is being referred to is the resting pressure. The American Heart Association has established a suggestion that the systolic blood pressure of an adult should be less than 140, and the diastolic blood pressure of an adult should be less than 90. This advice is in accordance with the American Heart Association's guidelines. It is possible that individuals who have blood pressure readings that fall within the range of 120-139/80-90 are at risk for developing hypertension. Because hypertension is a condition that may be induced by high blood pressure, this is the reason why this is the case. It is generally advised that these people make alterations to the dietary habits and lifestyle choices that they presently engage in in order to lessen the possibility that they may develop hypertension in the future. Even if they are inside

this range, there are some people who do not need medicine at this time while others fall within this range. The folks in question do not have a need for medicine. People who have stage I hypertension (140-159/90-99) or stage II hypertension (160 or higher above 100 or higher) may need medication for blood pressures as low as 120/70, as stated by Juan (2013) and Susan (2010). Stage I hypertension is when the blood pressure is between 140 and 159/90-99. In spite of the fact that their blood pressure is lower than that, this occurs nevertheless. Stage I hypertension, which is defined as 140-159/90-99, is the most common kind of hypertension. It is also the most common type of hypertension. The severity of the individual's ailment is taken into account in order to determine whether or not this is the case based on the evaluation. In particular, this is the case for individuals who possess other features that, when considered as a whole, increase the likelihood that they may acquire cardiovascular disease.

Literature Review

Types and Description of blood pressure

Type of hypertension	Description	Blood pressure range
Essential hypertension (major type)	Chronic elevation in blood pressure with no underlying disease	Both systolic and diastolic blood pressure are elevated more than 140/ 90 mmHg
Secondary hypertension (second common type)	Chronic elevation in blood pressure due to underlying pathology (mostly due to renal problems)	Both systolic and diastolic blood pressure are elevated more than 140/ 90 mmHg
Malignant hypertension	When blood pressure is severely elevated and causes an organ damage	Both systolic and diastolic blood pressure are elevated but the diagnosis made mainly when the diastolic blood pressure is higher than 130 mmHg
Isolated systolic hypertension	Common in elderly due to the loss of elasticity of major blood arteries	The systolic blood pressure is higher than 140 mmHg while the diastolic blood pressure is close to the normal range
Resistant hypertension	When more than three different antihypertensive agents are prescribed and blood pressure is still elevated	Both systolic and diastolic blood pressure are elevated more than 140/90 mmHg

Causes of Hypertension: Understanding Primary and Secondary Factors

Primary (essential) hypertension and secondary hypertension are the two forms of hypertension that may be distinguished from one another. Hypertension, often known as high blood pressure, is a frequent term. 95% of people have primary hypertension, also known as essential hypertension. This form of hypertension accounts for about 95% of all occurrences of hypertension and often manifests itself between the ages of 20 and 50. There is a possibility that it is impacted by genetic factors, as shown by a favorable ancestral past. In essential hypertension, the following symptoms are present:

- No evidence of secondary causes has been found.
- Stability for an extended length of time if it is not regulated.
- The possibility of a misdiagnosis as other forms of hypertension, such as white-coat hypertension (in which the patient's blood pressure is high in a medical setting but normal at home) or masked

hypertension (in which the patient's blood pressure is normal in a medical setting but higher in other surroundings).

• The presence of occasional discrepancies, such as normal readings at home or during ambulatory monitoring, but elevated values in clinical settings, is one example of this.

- There are five percent of secondary causes of hypertension. Secondary hypertension, which accounts for around five percent of all instances, often manifests itself under certain circumstances or is associated with other health problems. On average, it manifests itself either before the age of 20 or beyond the age of 50, and in contrast to primary hypertension, it does not frequently include a family history. Among the features are:
- The presence of hypertensive urgency or significant elevations in blood pressure that call for quick medical treatment.
- Resistance to the traditional therapies for hypertension is a concern.

- The usage of certain drugs or situations and the following are examples of possible triggers:

Exogenous Substances: Things like corticosteroids, medications used for oral contraception, cocaine, and black licorice are examples.

Renal Issues: Acute kidney injury (AKI), chronic kidney disease (CKD), glomerulonephritis, and renovascular disease (including unilateral and bilateral renal artery stenosis) are all included in this category.

Mechanical Problems: Including but not limited to obstructive sleep apnea and aortic coarctation.

Endocrine Disorders: Glucocorticoid excess (Cushing syndrome or illness), catecholamine excess (pheochromocytoma), mineralocorticoid excess (primary aldosteronism), thyroid function abnormalities (both hyperthyroidism and hypothyroidism impact blood pressure differently), and hyperparathyroidism are some of the conditions that are included in this category.

Pregnancy: In particular, problems such as hypertension seen during pregnancy.

The underlying causes of hypertension need to be addressed in order to properly control the disease (Calgary, 2020). Therefore, it is essential to have a thorough understanding of these differences in order to effectively manage and treat hypertension efficiently.

2.3 Symptums of hypertension

Symptoms	
Very high blood pressure	Increased sweating
Chest pains	Tinnitus – ringing or buzzing in ears
Headache	Heart palpitations
Fatigue	General feeling of unwellness
Shortness of breath	Increased urination frequency
Convulsion	Blurred vision
Dizziness	Double vision
Nausea	Flushed face
Vomiting	Nose bleeds
Anxiety	Pale skin
Changes in vision	Red skin

Figure (1) Common symptoms of hypertension. (Sarwar,2013).

Pathophysiology of Hypertension

High blood pressure, commonly known as hypertension, is a chronic condition that is defined by elevated blood pressure over a prolonged period of time. Hypertension is also known as high blood pressure. One of the possible consequences of this illness is an increased risk of morbidity and mortality, in addition to the potential for significant organ damage. Blood pressure is the real measurement of blood pressure, and it is obtained by multiplying the output of the heart by the resistance of the systemic vascular system. Patients who are diagnosed with arterial hypertension may have an increase in either their cardiac output or their systemic vascular resistance, or perhaps a combination of both of these cardiovascular diseases. When compared to older adults, younger people often have a greater cardiac output. On the other hand, older folks are more likely to have increased systemic vascular resistance and vascular stiffness. According to Messerli et al. (2011), this phenomenon may be principally attributable to alterations that include the activation of α-adrenoceptors and the production of vasoconstrictive peptides, such as angiotensin and endothelins. The vasoconstriction that takes place as a result of these adjustments is brought about by an increase in the amount of cytosolic calcium that is present in the smooth muscle around the blood vessels. Furthermore, growth hormones like angiotensin and endothelins are responsible for encouraging vascular remodeling, which eventually leads to an increase in the mass of vascular smooth muscle. This is the final outcome of the process. This remodeling, in combination with an increase in the systemic vascular resistance, leads the left ventricle to be exposed to extra strain, which eventually culminates in left ventricular hypertrophy and diastolic dysfunction on the left side of the heart. When it comes to younger individuals, the pressure that is produced by the left ventricle of the heart is often not great. The majority of the waves that are reflected by the peripheral vasculature occur after systole, which contributes to an improvement in coronary perfusion. McKenzie et al. (2009) state that as people get older, they experience an increase in arterial stiffness, which in turn causes an increase in pulse pressure, which is a significant indicator of coronary heart disease. Alterations in the timing of the pulse, from early diastole to late systole, result in an increase in the afterload generated by the left ventricle, which further contributes to

hypertrophy. This is because the left ventricle is responsible for producing the afterload. Individuals who have hypertension often have elevated levels of norepinephrine release and sensitivity, in addition to heightened responses to stress. The autonomic nervous system is responsible for a significant portion of the regulation of blood pressure, which is the reason for this. Additionally, there is a reduction in the sensitivity of baroreceptors, as well as a recalibration of the baroreflexes to the new levels. Both of these changes occur simultaneously. The renin-angiotensin system is another significant contributor to hypertension. This system is particularly suppressed in circumstances such as primary hyperaldosteronism, which is a condition that causes hypertension. One of the types of hypertension that the system is most often connected with is referred to as renovascular hypertension. individuals may exhibit either low-renin or high-renin hypertension, with the latter being often linked to greater risks of myocardial infarction and related cardiovascular disorders. There is a chance that individuals may exhibit both types of hypertension.
One of the contributing factors to hypertension is pressure natriuresis, which refers to the incorrect management of blood volume and the connection between blood pressure and salt excretion. Pressure natriuresis is an important component of hypertension. This relationship is altered in essential hypertension, which often presents itself as either a shift to higher pressures with salt-insensitive hypertension or a decreased reactivity with salt-sensitive hypertension (Wald et al., 2009). Both of these manifestations are common types of hypertension. Both of these manifestations are instances that are typical of the symptoms associated with essential hypertension. As a result of the fact that the resetting of pressure natriuresis is a crucial component in the development of hypertension, it is suggested that there is a complex interaction between the systemic components that contribute to this clinical condition.

TARGET ORGAN "BRAIN":

Over the course of the last four decades, the incidence of stroke has decreased by more than forty percent in nations with high incomes. Nevertheless, over the same time period, the number of strokes that occur in nations with low and moderate incomes has increased by a factor of two.

1. A prominent element that contributes to the increasing risk of stroke is the maturation of the global population, which has been recognized as a significant contributor. If we consider that age is one of the most significant risk factors for stroke, then the increasing number of senior people indicates that there is a greater percentage of people who are at risk.

2. The reduction in the number of strokes that occur in nations with high incomes may be ascribed to the improvement in the treatment of cardiovascular risk. Ischemic strokes, which are caused by thrombosis or embolism, account for roughly 80 percent of strokes that occur in Western nations. Hemorrhagic strokes account for the remaining 20 percent. The ability to differentiate between hemorrhagic and ischemic strokes is essential for making choices on stroke management and patients' treatment.

Thrombosis and embolism are the primary factors that contribute to the development of an ischemic stroke. One of the most prevalent underlying conditions that leads to thrombotic ischemic stroke is atherosclerosis, which is characterized by the rupture of plaque. The presence of conditions such as high-grade stenosis of the internal carotid artery, fibromuscular dysplasia (FMD), arteritis (including giant cell and Takayasu), and vascular dissection might also be factors that lead to the development of thrombotic ischemic stroke. On the other hand, embolic stroke is often brought on by embolization from a variety of causes, with atrial fibrillation being the most prevalent cause of embolization. According to Mauricio and Gisele (2004), the presence of

several infarct sites in distinct arterial beds is indicative of the fact that the embolism started from the heart, and perhaps the aorta as well.

TARGET ORGAN "HEART":

Left Ventricular Hypertrophy (LVH):

An adaptive process that happens in response to the pressure load that is exerted on the heart is the initial cause of left ventricular hypertrophy (LVH). Its purpose is to lessen the stress on the wall of the heart while also preserving the function of the left ventricle (LV) pump and the ejection fraction. As a consequence, the wall thickness grows while the internal diameter of the left ventricle (LV) shrinks, which leads to a concentric pattern of left ventricular hypertrophy (LVH). The processes of left ventricular remodeling develop over time, which finally results in left ventricular dilatation, also known as eccentric left ventricular hypertrophy (LVH). These processes include thickening and shortening of hypertrophied muscle fibers, increased collagen content in the perivascular and interstitial areas, and ultimately, LV dilation. One thing that should be kept in mind, however, is that blood pressure (BP) alone can only account for a tiny percentage of the fluctuation in left ventricular mass (LVM). A number of non-hemodynamic variables, including body mass index, dietary salt consumption, hereditary factors, and activation of the sympathetic nervous system and renin-angiotensin system (RAS), are additional contributors to the development of low-volume cardiovascular disease (LVM) and the degree to which it manifests itself. One may evaluate left ventricular hypertrophy (LVH) in hypertension using a variety of techniques, each of which has a unique level of sensitivity and specificity. LVH, as measured by electrocardiogram (ECG) or echocardiogram, has been shown to be an independent risk factor for unfavorable cardiovascular and all-cause mortality outcomes (Pickering and White, 2008). This has been proven by epidemiological studies such as the one described above. An interesting observation is that left ventricular hypertrophy (LVH) detected by electrocardiogram (ECG) and echocardiogram (ECHO) shows separate pathogenetic elements connected to the adaptive process of LVH. According to Hsia et al. (2007), both electrocardiograms and echocardiographic left ventricular hypertrophy are independently related with death.

Heart Failure:

According to the findings of epidemiological research, hypertension is the most prevalent underlying cause of congestive heart failure (CHF). According to James et al. (2017), many antihypertensive medications, including ACE inhibitors, ARBs, aldosterone antagonists, beta-blockers, diuretics, and angiotensin receptor neprilysin inhibitors, are now considered to be standard care for the management of congestive heart failure (CHF). These medications not only reduce afterload, but they also counteract the neuroendocrine stimulation that is inherent in CHF.

Heart Failure with Preserved Ejection Fraction (HFpEF):

Alterations in left ventricular relaxation and filling patterns are the primary characteristics of diastolic dysfunction in hypertension. This dysfunction may occur prior to the onset of systolic dysfunction. Despite the fact that the ejection fraction (HFpEF) is preserved, clinical symptoms of heart failure may be brought on by diastolic dysfunction, which is often linked with left ventricular remodeling and concentric left ventricular hypertrophy. In order to diagnose heart failure with preserved ejection fraction (HFpEF), it is necessary to rule out various noncardiac causes of heart failure-like symptoms (Hsia et al., 2007). This task might be difficult.

TARGET ORGAN "KIDNEY":

Glomerular Damage:


Alterations that predominantly impact the tiny vessels have the potential to cause harm to the glomeruli. The constriction of the afferent arterioles that occurs in hypertension leads to partial ischemia of the glomerular tuft, which in turn leads to a decrease in filtration. The kidney is able to function for a longer amount of time as a result of hyperfiltration and hypertrophy of the remaining nephrons. This is the reason why end-stage renal disease (ESRD) is uncommon in hypertension people who do not have any additional risk factors, such as obesity, smoking, or diabetes mellitus. The condition known as hyaline arteriolosclerosis occurs when hyalinosis, which is defined by the deposition of hyaline material, primarily affects the arterioles that are responsible for nerve impulse transmission. Glomerulosclerosis, on the other hand, is the term used to describe the condition that occurs when the glomerular capillaries are affected. Glomerulosclerosis may be identified as focal if it affects just a subset of glomeruli, or as diffuse if it affects a greater percentage of glomeruli. Both of these classifications are possible. According to Mulvany and Aalkjaer (1990), diffuse glomerulosclerosis is often linked to nephrotic syndrome as well as increasing renal impairment.

Renal Fibrosis:

Renal fibrosis is a clinical characteristic that is often seen in patients with chronic kidney disease (CKD). It is distinguished by the abnormal buildup of extracellular matrix components in the interstitium of the kidneys. It is regarded as the ultimate common route that leads to renal illness that has reached the terminal stage. Through a variety of different processes, hypertension may play a role in the development of renal fibrosis. These mechanisms include the stimulation of the renin-angiotensin-aldosterone system, oxidative stress, inflammation, and endothelial dysfunction. In accordance with Liu (2011), these mechanisms encourage the proliferation of fibroblasts and myofibroblasts, which ultimately leads to the deposition of collagen and other proteins that are found in the extracellular matrix.

Renal Dysfunction:

One of the potential outcomes of chronic hypertension is the progression of renal impairment, which may eventually lead to end-stage renal disease (ESRD). A steady reduction in the glomerular filtration rate (GFR) and the development of proteinuria are two characteristics that are associated with the loss in renal function. Renal failure that is caused by hypertension is often linked to structural changes in the kidney. These alterations include glomerular damage, tubulointerstitial fibrosis, and vascular abnormalities. According to James et al. (2017), poor renal function may further worsen hypertension by activating a variety of processes. These mechanisms include the retention of salt and water, an increase in the activity of the sympathetic nervous system, and a dysregulation of the renin-angiotensin-aldosterone system (RAAS).



The haemodynamic forces that are present in a blood vessel are shown in Figure 1. According to Wright et al. (2015), distension pressure causes a radial force to be applied to the vessel wall, which in turn causes the wall to get longer and the vascular cells to align themselves in a direction that is perpendicular to the force that is being applied. Through the friction of the blood against the vessel wall, shear stress is formed. This stress is applied in a direction that is parallel to the surface of the vessel. The endothelium lining of the blood artery is the one that is directly responsible for bearing the effects of shear stress. As stated by Giles et al. (2015), the shear stress (τ) is directly related to the blood viscosity (μ) and the rate of flow (Q). The equation for τ is as follows: $\tau = 4\mu Q/\pi r3$, where r represents the vessel radius.

Hypertension and Cognition:-

The majority of the research that has been done on hypertension suggests that it is associated with cognitive impairment; nevertheless, there are studies that have produced contradictory results, which imply that there is a U-shaped association between blood pressure and cognitive performance. It is important to note that inferior cognitive function has been shown to be associated with both low and extremely high blood pressure levels. It is essential to keep the cerebral perfusion pressure at a suitable level in order to ensure normal brain perfusion, which requires blood pressure levels that are adequately high (Juraschech et al., 2021). There is a correlation between sustained increased blood pressure and rapid cognitive deterioration, according to the majority of research that follow participants over time. persons who have hypertension often have a worse level of performance in a variety of cognitive domains, such as learning, memory, attention, visuospatial skills, psychomotor abilities, and executive functioning, in comparison to persons who have normal blood pressure.

Evidence suggests that reducing blood pressure may greatly decrease the risk of dementia and cognitive decline in older age groups. This is especially true for those who are in cognitive decline. The administration of treatment for hypertension has been shown to mitigate the deterioration in scores on the mini-mental state test and to reduce the occurrence of dementia in comparison to the absence of treatment. It is probable that poorly regulated hypertension will raise the risk of cognitive impairment, including vascular dementia and

other kinds of dementia. This is the case despite the fact that the link between blood pressure and cognitive performance is difficult to understand. Therefore, it is vital to provide proper patient education about the possible hazards associated with hypertension. This is particularly true when taking into consideration the fact that hypertension is asymptomatic, which may discourage patients from seeking treatment (Kaelber et al., 2016; Skoog I, 1998; Spence JD, 1996).

DIAGNOSIS OF HYPERTENSION

Traditional methods of measuring blood pressure require the use of a cuff that is worn on the upper arm. On the other hand, recent studies have shown that it may be difficult to reliably measure blood pressure in individuals who have bigger upper arms, particularly those who have a body mass index (BMI) that is more than 35 kg/m2 (Irving, 2016). Concerns have been raised about the phenomenon known as "hidden under-cuffing," which occurs when the cuff bladder is too tiny for the size of the arm, which may result in blood pressure readings that are erroneously exaggerated. Comparing the diagnostic accuracy of blood pressure measures taken from the forearm, wrist, and fingertip to that of upper arm cuffs that fit appropriately in obese patients was the subject of a systematic review and meta-analysis that was carried out in the year 2016. According to the results, wrist measures indicated the best diagnosis accuracy for hypertension when compared to upper arm cuffs (McManus, 2018). This was the case when the wrist was held at the same level as the heart.

Blood pressure measurements fall into several categories

- **Normal blood pressure**: Blood pressure is considered normal if it's below 120/80 mm Hg.
- **Elevated blood pressure**: Elevated blood pressure refers to a systolic pressure ranging from 120 to 129 mm Hg and a diastolic pressure below 80 mm Hg (not above). This condition is also known as prehypertension.
- **Stage 1 hypertension**: Stage 1 hypertension is characterized by a systolic pressure ranging from 130 to 139 mm Hg or a diastolic pressure ranging from 80 to 89 mm Hg.
- **Stage 2 hypertension**: Stage 2 hypertension indicates more severe hypertension, with a systolic pressure of 140 mm Hg or higher or a diastolic pressure of 90 mm Hg or higher.
- Blood pressure measurement higher than 180/120 mm Hg constitutes a medical emergency that requires urgent care. Since blood pressure can fluctuate throughout the day and may rise during medical appointments (known as white coat hypertension), doctors typically conduct multiple blood pressure readings at three or more separate appointments before diagnosing high blood pressure (Doshi, 2010).

Prevention of hypertension

There is evidence that the incidence of cardiovascular disease and hypertension is quickly growing in sub-Saharan Africa (Nissinen A1988). Just under twenty percent of hypertension respondents in Tanzania were aware of their diagnosis, around ten percent reported getting treatment, and fewer than one percent had their blood pressure under control (140/90 mmHg) (Edwards R- 2000). These findings appear from two surveys that

were conducted not too long ago. Twenty percent of South African Black males were aware of their hypertension, fourteen percent were receiving treatment, and only seven percent had their blood pressure under control (140/90 mmHg). On the other hand, forty-seven percent of South African Black females were aware of their hypertension, twenty-nine percent were receiving treatment, and fifteen percent had their blood pressure under control (140/90 mmHg) (Steyn, 2001).

Primary prevention of hypertension

By implementing suitable policies and educational programs, the goal is to either prevent or postpone the development of hypertension (Cicolini et al., 2014). This may be accomplished by reducing or modifying the risk factors of hypertension, such as the amount of salt consumed and being obese. Benefits may be produced over the whole range of blood pressure distribution as a consequence of changes in behavior at the population level (for example, a reduction in salt consumption or an increase in physical activity). Just a little drop in blood pressure may bring about a large reduction in the risk of cardiovascular disease. This is due to the fact that a sizeable proportion of individuals have blood pressure that is higher than the ideal range. According to Seedat (2000), the major method of preventing hypertension at the individual level is to form good lifestyle habits at a young age.

Pharmacological treatment Principles of drug management

- The fundamental advantage of antihypertensive medication is that it lowers blood pressure, which is one of its key advantages. In the absence of strong grounds, it is advised that patients with diabetes begin treatment with low-dose thiazide diuretics, such as hydrochlorothiazide at a dosage of 12.5 milligrams per day. This recommendation applies to a wide range of patient groups. This method has been shown to be effective as the foundation of therapy in a number of outcome studies that have been conducted with hypertensive patients (Hermida et al., 2020; Seedat, 1999).
- • It is permissible to either raise the dosage of the same medicine (if side effects are not present) or add a minor dose of a second drug from a different class if a low dose of a single drug is well tolerated but fails to sufficiently regulate blood pressure. This is the case if additional drugs from a different class are also added. Whenever possible, generic formulations of any and all medications should be prescribed. Thiazide-like diuretics, long-acting calcium channel blockers (CCBs), beta-blockers, angiotensin-converting enzyme inhibitors (ACEIs), and reserpine are some of the common choices available. According to Staessen et al. (2003), Hypertens (2003), JAMA (2003), and Douglas (2003), the majority of patients need two or more antihypertensive medications in order to experience the desired effects of therapy. Diuretics are often prescribed as one of these medications.
- Methyldopa, clonidine, alpha-blockers like prazosin and doxazosin, and hydralazine are examples of pharmaceuticals that have a centrally acting mechanism of action. These medications should be kept for instances of hypertension that is resistant to treatment.
- • Because there have not been any mega-trials that directly compare angiotensin II receptor antagonists (ARAs) with angiotensin-converting enzyme inhibitors (ACEIs), generic ACEIs are favored over ARAs. On the other hand, there are circumstances in which ACEIs are not well tolerated, such as when a cough or angioedema affects the patient.

- In order to improve adherence to treatment and reduce the amount of variation in blood pressure, it is strongly suggested that patients take once-daily long-acting medications that have an effectiveness that lasts for twenty-four hours. In addition, this strategy may provide a higher level of protection against significant cardiovascular events and the development of damage to the organs that are the focus of the treatment.
- In Sub-Saharan Africa (SSA), it is observed that loop diuretics such as furosemide and diazepam are used rather often during treatment. On the other hand, diazepam is ineffectual, and the use of loop diuretics over an extended period of time may entail hazards. In some circumstances, such as congestive heart failure, severe renal disease, or hypertension that is resistant to treatment, furosemide should be retained for use for treatment.
- Patients with hypertension who are black often have a poor response to beta-blockers, ACE inhibitors, or ARAs, unless they are taken with a thiazide diuretic or similar medication. CCBs, vasodilators, and thiazide-like diuretics have been proven to be more beneficial for this demographic, according to research conducted by Bhatt et al. (2014).
- Although beginning treatment with two medications, even at a modest dosage, may put patients in contact with agents that are not essential, the benefits of doing so far exceed the drawbacks. According to Aziz et al. (2019), these benefits include better efficacy as a result of multiple modes of action, improved tolerance with fewer side effects, and enhanced adherence via the use of fixed low-dose combinations contained inside a single tablet.
- Several other combinations of two medications have been proven to be effective and well tolerated. These combinations include thiazide diuretic and beta-blocker, thiazide diuretic and ACEI or ARA, CCB (dihydropyridine) and beta-blocker, CCB and ACEI or ARA, CCB and thiazide diuretic, and alpha-blocker and beta-blocker. In the event that it is required, further combinations, such as those that include central agents or those that involve ACEI and ARA, may be used (Seedat, 2002).
- The monitoring of the patient's reaction to treatment is a vital component of the follow-up therapy that is required. It is recommended that patients be visited at regular intervals, such as once every one to two months, throughout the stabilization phase. This should continue until the patient's blood pressure levels are under satisfactory control. The major objective of healthcare practitioners is to manage additional risk factors such as systolic and diastolic blood pressure while simultaneously maintaining the goal levels of blood pressure. It is advised that blood pressure be lowered gradually and carefully in order to reduce the risk of adverse effects and problems, promote compliance, and guarantee that the treatment will be

Special situations Hypertension emergencies and urgencies:-

Situations that need immediate treatment of blood pressure in order to avoid or minimize damage to the organs that are the focus of the emergency are known as hypertensive crises. The treatment of these individuals may be administered on an outpatient basis; nevertheless, it is recommended that they be hospitalized in order to undergo a progressive and monitored decline in their blood pressure (Guan et al., 2020). When it comes to the majority of cases, high blood pressure should be controlled in the hospital with extreme caution and a gradual approach. In general, the objective is to produce a gradual reduction in blood pressure to a diastolic blood pressure of 100 mmHg as quickly as possible during the course of 48–72 hours. According to

research conducted by Cormick et al. in 2022, the brain arteries are only able to adjust to a reduction in blood pressure relatively slowly (over a few days or weeks), and a sudden drop in blood pressure may lead to a stroke or other issues.
There is no justification for a sudden drop in blood pressure in the event of a heart attack or stroke. High blood pressure is a typical occurrence after an ischemic stroke or primary intracerebral hemorrhage, and it is linked to a poor functional result. Despite this, the appropriate treatment of blood pressure in patients who have had an acute stroke is still uncertain.

The following drugs are often used.

(i) Long-acting calcium channel blockers are available. When using short-acting CCBs, sublingual or other forms should never be used.

(ii) ACE inhibitors, with modest dosages used initially. Before using the medicine, it is important to check the serum electrolytes for any signs of severe hyponatraemia or hyperkalemia.

(iii)-blockers

(iv)Diuretics. This may also potentiate the effects of the other classes mentioned above.

:**Hypertensive emergencies are life threatening. Common situations include:**

• Hypertensive encephalopathy.

• Cerebrovascular accident (CVA, stroke); CVA and stroke require a more prudent approach for the first 10 days. After stabilization, the blood pressure should be progressively decreased (Tikkanin et al,2015).

• Hypertensive heart failure presenting with acute left ventricular failure with severe pulmonary oedema (Tschanz et al,2020).

• Malignant hypertension.

• Dissecting aneurysm of the aorta.

• Eclampsia and pre-eclampsia.

• Unstable angina/myocardial infarction (J Hypertens 2003)

Conclusions and Recommendations:

1. The method of measuring blood pressure, which is a routine procedure in clinical practice, requires meticulous execution in order to function properly.

2. During training, there should be a strong focus placed on the common causes that result in inappropriate readings..

3. The practical features of blood pressure measurement have to be included in competency exams for all doctors and other medical professionals.

4. In addition to the implementation of a periodic maintenance schedule that is both cost-effective and efficient, it is essential to strictly prevent using equipment that is either faulty or inappropriate.

5.A customized maintenance routine need to be established in every healthcare environment where blood pressure readings are carried out.
6.Because mercury is a dangerous substance, handling mercury sphygmomanometers requires extreme attention, and there are systems in place to deal with any spills that may occur.
7.It is quite probable that mercury sphygmomanometers will be replaced as automation continues to improve; nonetheless, it is vital to validate the accuracy of electronically controlled devices independently.
8. It is important to pay attention to specialized methods for measuring blood pressure in different populations, such as youngsters, obese people, old people, and pregnant women.
9.For the purpose of future study, it should be a top priority to investigate the prevalence of hypertension and its effect inside our nation.

References:

1. Ahmed, A.F. , Johns . E.J .. 2012. The regulation of blood perfusion in the renal cortex and medulla by reactive oxygen species and nitric acid.
2. Archad, J.-M. , Fournier , A .. Mazouz , H. , Caride . V.J. , Penar . P.L ... Fernandez , L.A. , 2001. Protection against ischemia : a physiological function of the renin - angiotensin system . Biochem . Pharmacol . 261-272.
3. Azizi M, Sanghvi K, Saxena M, et al. Ultrasound renal denervation for hypertension resistant to a triple medication pill (RADIANCE-HTN TRIO): a randomised, multicentre, single-blind, sham-controlled trial. Lancet. 2021 Jun 26;397(10293):2476-86.
4. Azizi M, Schmieder RE, Mahfoud F, et al; RADIANCE-HTN Investigators. Six-month results of treatment-blinded medication titration for hypertension control following randomization to endovascular ultrasound renal denervation or a sham procedure in the RADIANCE-HTN SOLO Trial. Circulation. 2019 May 28;139(22):2542-53.
5. B. M. Egan, Y Zhao, and R. N. Axon, "US trends in prevalence, awareness, treatment, and control of hypertension, 1988–2008, Journal of the American Medical Association, vol . 303, no. 20, pp. 2043 2050, 2010.
6. Barquera S., I. Campos-Nonato, L. Hern andez-Barrera et al., "Hypertension in Mexican adults: results from the national health and nutrition sur vey 2006," Salud Publica de México, vol. 52, supplement 1, pp. S63-S71, 2010.
7. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood pressure.
8. Cicolini G, Simonetti V, Comparcini D, et al. Efficacy of a nurse-led email reminder program for cardiovascular prevention risk reduction in hypertensive patients: a randomized controlled trial. Int J Nurs Stud. 2014 Jun;51(6):833-43.
9. Cormick G, Ciapponi A, Cafferata ML, et al. Calcium supplementation for prevention of primary hypertension. Cochrane Database Syst Rev. 2022 Jan 11;1:CD010037.
10. Doshi H, Weder AB, Bard RL, Brook RD. Does "hidden undercuffing" occur among obese patients? Effect of arm sizes and other predictors of the difference between wrist and upper arm blood pressures. Journal of Clinical Hypertension. 2010;12:82-88.

11. Douglas JG, Bakris GL, Epstein M, Ferdinand KC, Ferrario C, Flack JM, et al. Management of high blood pressure in African Americans: consensus statement of the Hypertension in African Americans Working Group of the International Society on Hypertension in Blacks. Arch Intern Med 2003; 163:525–541
12. Duckitt K, Harrington D (2005) Risk factors for pre-eclampsia at antenatal booking: systematic review of controlled studies. BMJ 330(7491):565.
13. Duley L, Henderson-Smart D, Knight M et al. (2001) Antiplatelet drugs for prevention of pre-eclampsia and its consequences: systematic review. BMJ 322(7282):329–333.
14. Edwards R, Unwin N, Mugusi F, Whiting D, Rashid S, Kissima J, Aspray TJ, et al. Hypertension prevalence and care in an urban and rural area of Tanzania. J Hypertens 2000; 18:145–152.
15. ESH/ESC. 2003 European Society of Hypertension–European Society of Cardiology guidelines for the management of arterial hypertension. J Hypertens 2003; 21:1011–1053.
Ganong's Review of Medical Physiology Twenty-Third Edition
16. Giles TD, et al: Expanding the definition and classification of hypertension. J Clin Hypertens (Greenwich ISI Web of Science).
17. Guan Y, Dai P, Wang H. Effects of vitamin C supplementation on essential hypertension: A systematic review and meta-analysis. Medicine (Baltimore). 2020 Feb;99(8):e19274.
18. Gupta A, Perdomo S, Billinger S, et al. Treatment of hypertension reduces cognitive decline in older adults: a systematic review and meta-analysis. BMJ Open. 2020 Nov 17;10(11):e038971.
19. Hermida RC, Crespo JJ, Domínguez-Sardiña M, et al. Bedtime hypertension treatment improves cardiovascular risk reduction: the Hygia Chronotherapy Trial. Eur Heart J. 2020 Dec 21;41(48):4565-76.
20. Hsia J, Margolis KL, Eaton CB et al. (2007) Prehypertension and cardiovascular disease risk in the Women's Health Initiative. Circulation 115(7):855–860.
21. Irving G, Holden J, Stevens R, McManus RJ. Which cuff should I use? Indirect blood pressure measurement for the diagnosis of hypertension in patients with obesity: a diagnostic accuracy review. BMJ Open. 2016;6:doi:10.1136/bmjopen-2016-012429
22. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). JAMA. 2014 Feb 5;311(5):507-20.
23. Joint ESH/ESC Guidelines (2013) Guidelines for the management of arterial hypertension. Eur Heart J 34(28):2159–2219.
24. Juraschek SP, Hu JR, Cluett JL, et al. Effects of intensive blood pressure treatment on orthostatic hypotension: a systematic review and individual participant-based meta-analysis. Ann Intern Med. 2021 Jan;174(1):58-68.
25. Kaelber DC, Liu W, Ross M, Localio RA, Leon JB, et al (2016) Diagnosis and Medication Treatment of Pediatric Hypertension: A Retrospective Cohort Study for the Comparative Effectiveness Research Through Collaborative Electronic Reporting (CER2) Consortium. Pediatrics 138: 2016-2195
26. Mauricio Wajngarten and Gisele Sampaio Silva 2016. Hypertension and stroke : update on treatment.
27. McKenzie TJ, Lillegard JB, Young WF Jr. et al. (2009) Aldosteronomas – state of the art. Surg Clin North Am 89(5):1241–1253.
28. McManus RJ, Mant J, Franssen M, et al. Efficacy of self-monitored blood pressure, with or without telemonitoring, for titration of antihypertensive medication (TASMINH4): an unmasked randomised controlled trial. Lancet. 2018 Mar 10;391(10124):949-59.

29. Messerli F, Bangalore S, Makani H et al. (2011) Flash pulmonary oedema and bilateral renal artery stenosis: the Pickering syndrome. Eur Heart J 32(18):2231–2235.
30. Neal B, MacMahon S, Chapman N. Effects of ACE inhibitors, calcium antagonists and other blood pressure lowering drugs: results of prospectively designed overviews of randomized overviews of blood pressure.
31. Nissinen A, Bothig S, Granroth H, Lopez AD. Hypertension in developing countries. World Health Stat Q 1988; 41:141–154 oxide in the anaesthetised rat . Acta Physiol . (Oxf) 204 , 443 450
32. P. M. Kearney, M. Whelton, K. Reynolds, P. Muntner, P K. Whelton, and J. He, "Global burden of hypertension: Analysis of worldwide data," The Lancet, vol. 365, no. 9455, pp. 217-223, 2005.
33. Pickering TG, White WB (2008) Ash Position Paper: Home and ambulatory blood pressure monitoring. When and how to use self (home) and ambulatory blood pressure monitoring. J Clin Hypertens (Greenwich) 10(11):850–855.
34. Rabi DM, McBrien KA, Sapir-Pichhadze R, et al. Hypertension Canada's 2020 comprehensive guidelines for the prevention, diagnosis, risk assessment, and treatment of hypertension in adults and children. Can J Cardiol. 2020 May;36(5):596-624.
35. Rana BK, Insel PA, Payne SH et al. (2007) Population-based sample reveals gene-gender interactions in blood pressure in white Americans. Hypertension 49(1):96–106.randomized trials. Blood Pressure Treatment Trialists' Colloboration. Lancet 2000; 356:1955–1964
36. Sarwar MS et al. Resistant Hypertension: Cause & Treatment … Drug Res 2013; 63: 217–223
37. Seedat YK. Hypertension in black South Africans. J Hum Hypertens 1999; 13:97–103
38. Seedat YK. Hypertension in developing nations in sub-Saharan Africa. J Hum Hypertension 2000; 14:739–747
39. Seedat YK. Malignant accelerated hypertension In: Mancia G, Chalmers J, Julius S, Saruta T, Weber M, Ferrari A, Wilkinson I (editors): Manual of hypertension. London: Churchill Livingstone Harcourt Publishers Limited; 2002, pp. 623–634.
40. Skoog I (1998) A review on blood pressure and ischaemic white matter lesions. Dement Geriatr Cogn Disord 9 Suppl 1: 13-19
41. Spence JD (1996) Cerebral consequences of hypertension: where do they lead? J Hypertens Suppl 14: S139-145. [Crossref]
42. Staessen JA, Wang JG and Birkenhager WH. Outcome beyond blood pressure control? Eur Heart J 2003; 24:504–514.
43. Steyn K, Gaziano TA, Bradshaw D, Laubscher R, Fourie J. South African Demographic and Health Coordinating Team. Hypertension in South African adults: results from the Demographic and Health Survey, 1998. J Hypertens 2001; 19:1717–1725.
44. Teresa M.Secciaa, Brasilina Carocciaa, and Lorenzo A. Calo`, 2017 Hypertensive nephropathy. Moving from classic to emerging pathogenetic mechanisms.
45. Tikkanen I, Narko K, Zeller C, et al; EMPA-REG BP Investigators. Empagliflozin reduces blood pressure in patients with type 2 diabetes and hypertension. Diabetes Care. 2015 Mar;38(3):420-8.
46. Tschanz CMP, Cushman WC, Harrell CTE, et al. Synopsis of the 2020 U.S. Department of Veterans Affairs/U.S. Department of Defense Clinical Practice Guideline: the diagnosis and management of hypertension in the primary care setting. Ann Intern Med. 2020 Dec 1;173(11):904-13.
47. Wald DS, Law M, Morris JK et al. (2009) Combination therapy versus monotherapy in reducing blood pressure: meta-analysis on 11,000 participants from 42 trials. Am J Med 122:290–300.

48. Walsh JB. Hypertensive retinopathy: description, classification, and prognosis. Ophthalmology 1982;89:1127-1131
49. WHO/ISH. 2003 World Health Organization/International Society of Hypertension Statement on Management of Hypertension. J Hypertens 2003; 21:1983–1992.
50. Wright JT Jr, Williamson JD, Whelton PK, et al; SPRINT Research Group. A randomized trial of intensive versus standard blood pressure control. N Engl J Med. 2015 Nov 26;373(22):2103-16.