MODERN MEDICINE AND PRACTICE
https://inovatus.es/index.php/ejmmp
EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 4 No. 9 (Sep - 2024) EJMMP ISSN: 2795-921X

EVALUATION OF THE MORPHOLOGICAL STATE OF THE PANCREAS IN DIABETES IN CHRONIC POISONING WITH PESTICIDES

Ubaydullayev Abbosxo'ja Xikmatilla o'g'li
Assistant of the department "Internal diseases, therapy in family medicine, military field therapy and hematology", Tashkent Medical Academy, Termiz branch

Saidov Jasur Baxtiyorovich
Teacher of the Department of "Medical Clinical Sciences", Termiz University of Economics and Service

Annotation: In this article, the opinions of our country's and foreign scientists are mentioned about the assessment of the morphological state of the pancreas in chronic poisoning with pesticides in diabetes mellitus. The evaluation of the morphological state of the pancreas in diabetes under chronic pesticide poisoning involves studying how long-term exposure to pesticides affects pancreatic tissue, particularly in individuals with diabetes. Chronic pesticide exposure has been shown to have detrimental effects on various organs, including the pancreas. In diabetes, the pancreas already undergoes significant morphological changes, particularly involving beta-cell dysfunction and loss. When combined with chronic poisoning by pesticides, the damage to the pancreas can be exacerbated, leading to altered endocrine and exocrine function.

Key words: Chronic hyperglycemia, hallmark of diabetes mellitus (DM), endocrine disruption, oxidative stress, inflammation, diabetes mellitus, hyaline degeneration, inflammatory infiltration.

Introduction.

Chronic hyperglycemia, a hallmark of diabetes mellitus (DM), is a metabolic disease that can be brought on by a number of variables, including genetic, environmental, and lifestyle choices. Among environmental variables, diabetes and associated pancreatic problems have been linked to long-term pesticide exposure. This paper examines the anatomical alterations that take place in the pancreas of diabetic patients, with a focus on persistent pesticide exposure.

The connection between diabetes and pesticides

Changes in pancreatic function and glucose metabolism can result from long-term pesticide exposure for the following reasons:

1. Endocrine Disruption: Some pesticides have the ability to interfere with insulin signalling pathways since they are endocrine disruptors.
2. Oxidative Stress: Pesticides have the ability to cause oxidative stress, which damages pancreatic beta cells' cells.[1]

[1] Adeola H. A., et al. "The Pancreatic Effect of Pesticides in Experimental Models." *Journal of Toxicology*, 2021.

3. Inflammation: Extended exposure can cause the pancreas to become chronically inflamed, which exacerbates insulin resistance.

Materials.

Diabetes Mellitus: Morphological Alterations in the Pancreas 1. Betacell Dysfunction:

Reduced Betacell Mass: Research indicates that necrosis or apoptosis have led to a decrease in the quantity of pancreatic islet cells.

Islet Hyperplasia or Hypertrophy: In Type 2 diabetes, compensatory processes may lead to hyperplasia at first, which is followed by beta cell burnout in the end.

2. Hyaline Degeneration: Amyloid deposits build up in the islets, especially in Type 2 diabetes, which causes betacell malfunction.[2]

3. Vascular Changes: This refers to the pancreatic stroma's increased fibrosis and vessel wall thickening, which alter blood flow and oxygenation.

4. Inflammatory Infiltration: increased in Type 1 diabetes, lymphocytic infiltration in the islets.

In Severe Pesticide Exposure

1. Acinar Cell Damage: Directly harmful pesticide impacts can cause necrotic alterations in acinar cells, as shown by histopathological testing.
2. Fatty Infiltration: Acinar cell steatosis may transpire, impairing the pancreas's ability to synthesise.
3. Inflammatory Response: The pancreas showed an increase in inflammatory cells, suggesting a persistent inflammatory process that may be related to pesticide exposure.[3]
4. Modified Islet Architecture: As a result of exposure to environmental pollutants, islet morphology was disrupted, resulting in increased fibrosis and changes in cellular composition.
5. Chronic Fibrosis: Chronic inflammation and damage cause fibrotic alterations in the pancreatic tissue.

The Joint Impact of Pesticide Exposure and Diabetes

Synergistic Damage: Prolonged poisoning can worsen the morphological alterations brought on by diabetes, which can result in more serious pancreatitis.[4]

Increased Risk of Diabetes Complications: People with diabetes are more likely to experience nephropathy and cardiovascular disorders as a result of their altered pancreatic shape.

Research and methods.

Methods of Diagnosis and Assessment

1. Histopathological Examination: Pancreatic tissue biopsy samples can shed light on cellular makeup and structural alterations.
2. Imaging Methods: MRI, CT scan, and ultrasound are helpful in determining the size, texture, and presence of any abnormal masses in the pancreas.

[2] Kaushik S., et al. "Impact of Environmental Toxins on Pancreatic Function in Diabetes." *Diabetes & Metabolic Syndrome Reviews*, 2020.
[3] Manna P., et al. "Pesticide-Induced Oxidative Stress and Pancreatic Dysfunction in Diabetes." *Toxicology Letters*, 2017.
[4] Mahajan R., et al. "Chronic Pesticide Exposure and Pancreatic Islet Dysfunction." *Environmental Health Perspectives*, 2019.

3. Biochemical Markers: Evaluation of oxidative stress levels, routine glucose metabolism tests, and inflammatory markers.[5]

Theoretical Underpinnings of Assessing the Morphological Status of the Pancreas in Diabetes Mellitus Patients Who Have Chronic Pesticide Poisoning

Because beta cells in the islets of Langerhans produce insulin, the pancreas is essential for the metabolism of glucose. Pesticide exposure over an extended period of time can cause morphological changes in pancreatic tissues, which may exacerbate diabetic mellitus (DM). To evaluate these alterations, a comprehensive theoretical methodology incorporating principles from pathology, endocrinology, and toxicology is necessary.

Results.

1. Toxicological Principles in the Theoretical Framework

The toxicity's mechanisms: It is essential to comprehend the cellular interactions of several pesticide types, such as glyphosate, carbamates, and organophosphates.

Pesticides may be lipophilic, which enables them to concentrate in fatty tissues like the pancreas. This is known as cellular uptake and distribution.

Biotransformation: The liver breaks down some pesticides into hazardous byproducts that can harm pancreatic cells.[6]

Relationships Between Dose and Response: The degree of morphological alteration in pancreatic tissues is frequently dose-dependent and can be measured by:

Finding the dosage at which appreciable changes in pancreatic morphology take place is known as the threshold level.

Cumulative Effects: Extended exposure may cause harm that worsens over time, requiring further research.

2. Diabetes's Pathophysiological Mechanisms

Betacell Dysfunction: The decrease of mass and function in the betacells causes reduced insulin production, which is a hallmark of diabetes. Chronic pesticide exposure can alter the following things:

Apoptosis of Beta Cells: Pesticides can cause oxidative stress, which causes insulin-producing cells to undergo programmed cell death.

Modified Insulin Signalling: Pesticides may interfere with insulin receptor signalling pathways, causing disruptions in glucose metabolism.

Increased Insulin Resistance: Toxins from the environment can make insulin resistance worse by:

Chronic inflammation has the potential to modify adipocyte function and elevate insulin resistance.

Endocrine Disruption: Pesticides have the ability to imitate or obstruct hormone signals, which exacerbates the insulin resistance linked to diabetes.

3. Evaluation of Histopathology

Morphometric Evaluation: The process of quantitatively evaluating pancreatic morphology includes:[7]

[5] Garcia-Arevalo M, et al. "Organophosphate pesticide exposure induces insulin resistance, disrupts glucose metabolism, and causes pancreatic beta-cell dysfunction." Toxicology Letters, 2014.
[6] King H, Rewers M. "Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults." Diabetes Care, 1993.

Mass of Islet Cells: Analysing Langerhans islets' quantity, size, and distribution can reveal information about the health of betacells.

Tissue Architecture: Fibrosis, fat infiltration, and necrosis are examples of structural alterations that may be seen by H&E staining.

Biochemical Indicators of Damage: Assessing the expression of particular proteins can offer further background information on morphological alterations:

Apoptosis markers: Bcl2 and Caspase 3 can show how well betacells are doing.

Oxidative Stress Markers: Oxidative damage may be measured by substances such as superoxide dismutase (SOD) levels and malondialdehyde (MDA).[8]

4. The Inflammatory Mechanisms

Cytokine Production: Extended exposure to pesticides can raise proinflammatory cytokines (such TNFα and IL6), which may contribute to pancreatic damage.

Particular Immune Response: Examining the amount of immune cells (such as lymphocytes and macrophages) in the pancreas might indicate how much inflammation is caused by pesticide exposure.

Discussion.

Methodologies for Assessment

1. Histological Methodologies

Light microscopy: Enables a thorough analysis of the morphology and cellular organisation of the pancreas.

Immunohistochemistry: May be used to evaluate the expression of certain proteins related to oxidative damage, inflammation, and apoptosis.

2. The study of electron microscopy

gives information on the ultrastructure of cells, making it possible to identify cellular changes that are not evident with light microscopy.

3. Assays for Biochemistry

Enzyme-Linked Immunosorbent Assay (ELISA): To measure oxidative stress indicators and inflammatory cytokines in pancreatic tissues.

Western Blotting: To measure apoptosis and stress response marker protein expression levels.

4. Methods of Molecular Biology

RTPCR: For examining the gene expression patterns in pancreatic tissues linked to apoptosis, inflammation, and oxidative stress.

Integrative Methodology

When evaluating the morphological status of the pancreas in diabetes mellitus caused by persistent pesticide exposure, the following details should be included:

Collaboration Across Disciplines: properly interpreting results by combining knowledge from endocrinology, pathology, and toxicity.

[7] Saxena P, et al. "Chronic pesticide exposure and pancreatic toxicity: Role of oxidative stress and inflammation." Journal of Endocrinology, 2010.
[8] Atkinson MA, Eisenbarth GS. "Type 1 diabetes: new perspectives on disease pathogenesis and treatment." Lancet, 2001.

Studies that track changes in morphology and function over time in order to determine a cause-and-effect link are known as longitudinal studies.

Models for Risk Assessment: using information from epidemiology and animal research to forecast human consequences.

Conclusion.

When examining the morphology of the pancreas in diabetics, particularly when persistent pesticide poisoning is present, notable pathological alterations are shown that may have a role in the development and exacerbation of the illness. It is essential to comprehend these changes in order to create preventative and treatment plans that work. It is imperative to do more study concentrating on the molecular routes of pesticide-induced damage to pancreatic tissues.

Recommendations.

Environmental Monitoring: To prevent chronic poisoning, regular evaluation of pesticide usage and exposure levels in agricultural settings is necessary.

Patient Education: Inform people who reside in farming areas about the possible health hazards that come with being exposed to pesticides.

Investigate the long-term impacts of pesticide exposure on the development of diabetes and pancreatic morphology using longitudinal research.[9]

In order to better understand and manage diabetes, this study emphasises the need of taking environmental factors like pesticide exposure into account. This will result in improved public health treatments and policies.

An integrated theoretical approach is required to evaluate the morphological status of the pancreas in the context of diabetes mellitus and chronic pesticide exposure. A thorough understanding of pancreatic health may be obtained by applying sophisticated histological and biochemical methods, comprehending the underlying toxicological concepts, and comprehending the pathophysiology mechanisms of diabetes. Having this basic understanding is essential to creating preventative and treatment plans that work for those who are at risk.

List of used literatures:

1. Garcia-Arevalo M, et al. "Organophosphate pesticide exposure induces insulin resistance, disrupts glucose metabolism, and causes pancreatic beta-cell dysfunction." Toxicology Letters, 2014.
2. King H, Rewers M. "Global estimates for prevalence of diabetes mellitus and impaired glucose tolerance in adults." Diabetes Care, 1993.
3. Saxena P, et al. "Chronic pesticide exposure and pancreatic toxicity: Role of oxidative stress and inflammation." Journal of Endocrinology, 2010.
4. Atkinson MA, Eisenbarth GS. "Type 1 diabetes: new perspectives on disease pathogenesis and treatment." Lancet, 2001.
5. Lenzen S. "The mechanisms of alloxan- and streptozotocin-induced diabetes." Diabetologia, 2008.
6. Joshi SC, Sharma P, Sharma N. Pesticides and pancreatic toxicity: A review. Indian J Pharmacol. 2011.
7. Al-Eryani L, Wahlang B, Falkner KC, et al. Pesticides induce pancreatic dysfunction: A role for environmental toxicants in diabetes. PLoS ONE. 2015.

[9] Lenzen S. "The mechanisms of alloxan- and streptozotocin-induced diabetes." Diabetologia, 2008.

8. Heindel JJ, Blumberg B, Cave M, et al. Metabolism disrupting chemicals and metabolic disorders. Reprod Toxicol. 2017.
9. Moreno JA, Peinado FM, Barbancho M, et al. Chronic exposure to organophosphates induces morphological changes in the pancreas. Toxicol Appl Pharmacol. 2018.
10. Hectors TL, Vanparys C, Van Der Ven K, et al. Environmental pollutants and type 2 diabetes: A review of mechanisms that can disrupt beta-cell function. Diabetologia. 2011.