MODERN MEDICINE AND PRACTICE
https://inovatus.es/index.php/ejmmp
EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 4 No. 9 (Sep - 2024) EJMMP ISSN: 2795-921X

ANEMIA AND ITS PREMATURE BIRTH COMPLICATION WITH CLINICAL SYMPTOMS

O'Rolova Sumbula Sobirovna Tibbyot
Assistant at Termiz Branch of Tashkent Medical Academy

Annotation: In this article, the opinions of our country's and foreign scientists about the causes of premature birth and anemia and its prevention are mentioned. Premature birth and anemia are significant health issues, particularly for pregnant women and newborns. Premature birth and anemia are both significant health concerns for pregnant women and newborns.

Key words: Premature birth, immature blood production, iron deficiency, blood loss during delivery, inadequate iron absorption, blood transfusions, increased risk of infections, early intervention.

Introduction.

Premature birth (born before 37 weeks of gestation) and anemia (low red blood cell count) are two significant health challenges that can impact a newborn's health and development. While they can occur independently, they are often intertwined, creating a complex situation with potential complications.

Here's a breakdown of their relationship:

Why Premature Birth Increases Risk of Anemia:

Immature Blood Production: Premature babies have immature bone marrow, which produces fewer red blood cells, leading to anemia.

Iron Deficiency: Premature infants often have low iron stores due to their shorter gestation period, making them susceptible to iron deficiency anemia.

Blood Loss During Delivery: Premature births may involve complications like placenta previa or placental abruption, resulting in blood loss for the infant.[1]

Inadequate Iron Absorption: Premature babies may have difficulty absorbing iron from breast milk or formula due to their underdeveloped digestive systems.

Blood Transfusions: Premature babies often receive blood transfusions due to various complications, which can dilute their own blood and contribute to anemia.

Impact of Anemia on Premature Infants:[2]

Delayed Growth and Development: Anemia can hinder growth and development due to reduced oxygen delivery to vital organs.

[1] Mayo Clinic Staff. (2023). Premature birth. Mayo Clinic.
[2] American Pregnancy Association. (2022). Premature labor and birth.

Materials.

Increased Risk of Infections: Anemia weakens the immune system, making premature infants more vulnerable to infections.

Brain Damage: Severe anemia can lead to brain damage due to oxygen deprivation.

Heart Problems: Anemia can increase workload on the heart, leading to heart problems.

Cognitive Impairment: Anemia in the early stages of life can impact cognitive development and learning abilities.

Addressing Anemia in Premature Infants:

Iron Supplementation: Iron supplementation is crucial for premature infants to prevent and treat iron deficiency anemia.

Blood Transfusions: In cases of severe anemia, blood transfusions may be necessary to increase red blood cell count.[3]

Early Intervention: Early detection and treatment of anemia are crucial to minimize potential complications.

Monitoring and Management: Regular blood tests and monitoring are vital to assess iron levels and ensure effective management of anemia.

Premature birth, also known as preterm birth, is one that occurs before 37 weeks of gestation. Babies born prematurely are at a higher risk of developing certain health complications, including anemia. Anemia is a condition characterized by a low level of red blood cells or hemoglobin in the blood, which can lead to symptoms such as fatigue, weakness, and pale skin.

Research and methods.

Premature babies are more susceptible to anemia for several reasons:

1. Reduced iron stores: Iron is essential for the production of red blood cells, and premature babies often have lower iron stores at birth compared to fullterm babies. This can predispose them to iron deficiency anemia.

2. Immature red blood cell production: Premature babies may have immature red blood cell production, leading to a decreased ability to produce an adequate number of red blood cells to transport oxygen efficiently.

3. Blood loss: Premature babies are at a higher risk of blood loss due to medical procedures like blood tests, blood transfusions, and frequent blood sampling.[4]

4. Nutritional deficiencies: Premature babies may have difficulty feeding and absorbing nutrients, including iron, which can contribute to anemia.

To prevent and manage anemia in premature babies, healthcare providers may recommend the following:

Iron supplementation: Premature babies may need iron supplements to support the production of red blood cells and prevent or treat iron deficiency anemia.

Monitoring: Healthcare providers will monitor the baby's hemoglobin levels and iron status through regular blood tests to detect and address anemia early.

[3] National Institutes of Health (NIH). (2020). Iron-deficiency anemia. NIH.
[4] Cleveland Clinic. (2023). Anemia. Cleveland Clinic.

Blood transfusions: In severe cases of anemia, premature babies may require blood transfusions to replenish their red blood cell levels.

Nutritional support: Adequate nutrition is essential for preventing anemia in premature babies. Healthcare providers may recommend fortified breast milk or formula to ensure the baby is receiving enough iron and other essential nutrients.

It is important for parents of premature babies to work closely with healthcare providers to monitor their baby's growth, development, and overall health, including the prevention and management of anemia. Early detection and intervention can help reduce the impact of anemia on the baby's health and development.[5]

Results.

Causes of Premature Birth and Anemia:

Premature Birth:

Medical Conditions:

Infections: Uterine infections, like chorioamnionitis, can trigger labor prematurely.

High Blood Pressure: Preeclampsia and other hypertensive disorders can damage the placenta and lead to premature birth.

Cervical Insufficiency: A weak cervix can lead to early dilation and premature birth.

Multiple Pregnancies: Carrying twins, triplets, or more increases the risk of premature birth.

Previous Premature Births: Women who have had premature births before are at increased risk of having another one.

Lifestyle Factors:

Smoking: Smoking during pregnancy can constrict blood vessels, hindering oxygen flow to the fetus and increasing the risk of premature birth.

Alcohol Consumption: Excessive alcohol use can harm the fetus and trigger premature labor.

Drug Use: Drug use during pregnancy, including illicit drugs and some prescription medications, increases the risk of premature birth.

Stress: Chronic stress can affect hormone levels and contribute to preterm labor.[6]

Other Factors:

Placenta Problems: Placental abruption or previa can disrupt oxygen and nutrient supply to the fetus, leading to premature birth.

Uterine Abnormalities: Structural abnormalities in the uterus can increase the risk of premature birth.

Genetic Factors: Some individuals may have genetic predisposition to premature birth.

Anemia:

Iron Deficiency: Insufficient iron intake or poor iron absorption can lead to iron deficiency anemia.

Nutritional Deficiencies: Deficiencies in other nutrients like vitamin B12 and folate can also contribute to anemia.

[5] March of Dimes. (2021). **What causes preterm labor and birth?** Retrieved from https://www.marchofdimes.org
[6] World Health Organization (WHO). (2020). **Preterm birth**. Retrieved from https://www.who.int

Chronic Blood Loss: Conditions like heavy menstrual bleeding, ulcers, or internal bleeding can lead to chronic blood loss and anemia.

Genetic Conditions: Some genetic disorders can affect red blood cell production, leading to anemia.

Chronic Diseases: Chronic diseases like kidney disease, cancer, or inflammatory bowel disease can cause anemia.[7]

Prevention Strategies:

Premature Birth:

Prenatal Care: Regular prenatal checkups allow for early detection and management of potential risk factors.

Healthy Lifestyle: Maintaining a healthy weight, quitting smoking, avoiding alcohol and drug use, and managing stress can significantly reduce the risk of premature birth.

Early Treatment of Medical Conditions: Prompt diagnosis and treatment of conditions like infections, high blood pressure, and cervical insufficiency are crucial.

Cervical Cerclage: A surgical procedure to strengthen the cervix can be performed to prevent premature birth in women with cervical insufficiency.

Anemia:

Balanced Diet: Consume a diet rich in ironrich foods like red meat, leafy green vegetables, beans, and lentils.

Iron Supplements: Discuss iron supplementation with your healthcare provider if you are at risk of or diagnosed with iron deficiency anemia.

Vitamin B12 and Folate Intake: Ensure adequate intake of these nutrients through diet or supplementation.

Regular Blood Tests: Get regular blood tests to monitor iron levels and identify anemia early.

Medical Management: Treat underlying medical conditions that may be contributing to anemia.

Discussion.

Premature birth and anemia in babies can have various causes, and it is essential to understand these factors to prevent and manage these conditions effectively. Here are some of the common causes and ways to prevent premature birth and anemia:

Causes of Premature Birth:

1. Multiple pregnancies: Women carrying twins or multiples are at a higher risk of delivering prematurely.

2. Infections: Infections in the mother, such as urinary tract infections or sexually transmitted infections, can increase the risk of premature birth.

3. Maternal health conditions: Certain conditions like high blood pressure, diabetes, or preeclampsia can lead to premature birth.

4. Previous premature birth: Women who have had a previous premature birth are at a higher risk of experiencing it again.[8]

[7] National Heart, Lung, and Blood Institute (NHLBI). (2021). **What is anemia?** Retrieved from https://www.nhlbi.nih.gov
[8] Centers for Disease Control and Prevention (CDC). (2022). **Iron-Deficiency Anemia**. Retrieved from https://www.cdc.gov

5. Lifestyle factors: Smoking, drug use, or heavy drinking during pregnancy can increase the chances of premature birth.

Prevention of Premature Birth:

1. Prenatal care: Regular prenatal checkups are crucial for monitoring the health of both the mother and the baby and addressing any potential risk factors.

2. Healthy lifestyle: Maintaining a healthy diet, staying physically active, and avoiding harmful substances like tobacco and alcohol can help prevent premature birth.

3. Managing chronic conditions: Proper management of chronic conditions like diabetes or high blood pressure can reduce the risk of premature birth.

4. Stress management: Stress can impact pregnancy outcomes, so it's important for expectant mothers to manage stress levels through relaxation techniques or counseling.

Causes of Anemia in Babies:

1. Iron deficiency: Babies may develop anemia due to insufficient iron stores, which can occur if they are not getting enough ironrich foods.

2. Premature birth: Premature babies have lower iron stores and are at a higher risk of developing anemia.

3. Blood loss: Babies may experience blood loss due to medical procedures, such as blood tests or surgeries, leading to anemia.

4. Inadequate nutrition: Babies who are not getting enough ironrich foods or proper nutrition may develop anemia.

Prevention of Anemia in Babies:

1. Ironrich diet: Breastfed babies should receive ironfortified formula or iron supplements if needed. Once solid foods are introduced, ironrich foods like fortified cereals, meats, and green leafy vegetables should be included in the diet.

2. Prenatal care: Mothers should receive adequate prenatal care to ensure they have optimal iron levels during pregnancy, which can prevent anemia in babies.

3. Iron supplementation: Premature babies or babies born with low iron stores may require iron supplements to prevent or treat anemia.

4. Monitor for symptoms: Parents should be aware of the signs of anemia, such as pale skin, fatigue, or weakness, and seek medical attention if they observe any concerning symptoms.

Conclusion.

Premature birth significantly increases the risk of anemia in newborns. This complex interplay can negatively impact the infant's health and development. Early intervention, proper management, and monitoring are essential to ensure optimal health outcomes for premature infants suffering from anemia.

Preventing premature birth and anemia requires a multifaceted approach that addresses both medical and lifestyle factors. Early detection, proper management, and preventive measures are crucial to ensure a healthy pregnancy and a healthy start for newborns.

By understanding the causes of premature birth and anemia in babies and taking preventative measures, parents and healthcare providers can work together to promote the health and wellbeing of the newborn. Early detection and intervention are key in preventing and managing these conditions effectively.

List of used literatures:

1. Mayo Clinic Staff. (2023). Premature birth. Mayo Clinic.
2. American Pregnancy Association. (2022). Premature labor and birth.
3. National Institutes of Health (NIH). (2020). Iron-deficiency anemia. NIH.
4. Cleveland Clinic. (2023). Anemia. Cleveland Clinic.
5. March of Dimes. (2021). What causes preterm labor and birth? Retrieved from https://www.marchofdimes.org
6. World Health Organization (WHO). (2020). Preterm birth. Retrieved from https://www.who.int
7. National Heart, Lung, and Blood Institute (NHLBI). (2021). What is anemia? Retrieved from https://www.nhlbi.nih.gov
8. Centers for Disease Control and Prevention (CDC). (2022). Iron-Deficiency Anemia. Retrieved from https://www.cdc.gov