MODERN MEDICINE AND PRACTICE
https://inovatus.es/index.php/ejmmp
EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 5 No. 12 (Dec - 2025) EJMMP ISSN: 2795-921X

THE EFFECTIVENESS OF PROBING THE LACRIMAL DUCTS IN CONGENITAL DACRYOCYSTITIS

Boboev Saidavzal Abdurakhmonovich
Department of Ophthalmology, Samarkand State Medical University

Abdurakhmonova Sitora Saidavzalzoda
Samarkand region cardiology, dispensary doctor ophthalmologist

Abstract: Probing of the lacrimal drainage system is regarded as the primary interventional method for congenital dacryocystitis when conservative management fails. This procedure aims to eliminate obstruction at the valve of Hasner, restore physiological tear flow, and prevent recurrent infection. Clinical evaluation demonstrates that timely probing results in rapid resolution of epiphora, reduction of mucopurulent discharge, and prevention of chronic inflammation. Success rates depend on age, obstruction severity, anatomical variations, and operator experience. This analysis provides a detailed assessment of probing effectiveness, demonstrating that early intervention ensures improved functional recovery, minimizes long-term complications, and reduces the need for secondary surgical procedures. Probing of the nasolacrimal pathway is a key intervention for persistent congenital dacryocystitis, providing mechanical relief of distal blockage and enabling physiologic tear passage. In infants who do not respond to massage and antimicrobial therapy, this minimally invasive technique ensures rapid improvement in tearing, discharge, and inflammatory symptoms. Its effectiveness depends on age, type of obstruction, and structural flexibility of the duct. Early application consistently demonstrates superior recovery, fewer recurrences, and minimal complications. Expanded clinical evaluation shows that probing not only restores patency but also prevents chronic disease evolution that may lead to recurrent infections, cutaneous fistula formation, or sac distension.

Keywords: congenital dacryocystitis, nasolacrimal duct obstruction, probing, lacrimal drainage system, neonatal ophthalmology, tear flow restoration.

Introduction:

Congenital dacryocystitis represents a prevalent neonatal condition characterized by persistent tearing, mucous accumulation, and recurrent infections resulting from failure of canalization at the distal nasolacrimal duct. Although initial management typically involves massage of the lacrimal sac and topical antimicrobial therapy, a significant proportion of infants require surgical intervention to resolve obstruction and restore drainage. Probing of the nasolacrimal duct offers a minimally invasive yet highly effective method for treating persistent obstruction, particularly when performed within the age range most favorable for anatomical responsiveness. Understanding the mechanisms of obstruction resolution, ideal timing for intervention, and predictive factors for successful probing is essential for optimizing therapy and preventing chronic disease progression, which may lead to recurrent dacryocystitis, abscess formation, and compromised ocular surface health. Congenital dacryocystitis develops from incomplete canalization of the

lower nasolacrimal system, resulting in stagnant tear accumulation and subsequent bacterial proliferation. Infants typically present with persistent watering, mucous reflux on pressure, eyelid irritation, and recurrent episodes of inflammation. Initial conservative therapy targets membrane dissolution and infection control; however, when obstruction remains, interventional management becomes essential. Probing provides a structured approach by mechanically opening the distal barrier, allowing tears to drain freely and preventing further complications. Determining the optimal timing and understanding factors influencing outcomes are central to maximizing therapeutic success. Early decision-making plays a crucial role in reducing long-standing obstruction, limiting fibrosis, and avoiding future surgical procedures. Assessing the clinical value of this technique is therefore important in ensuring effective neonatal ophthalmologic care.

Research Methods and Approaches:

A systematic assessment was conducted using clinical charts, prospective observational data, and published studies focused on congenital nasolacrimal duct obstruction treated with probing between 2012 and 2024. Search parameters included “congenital dacryocystitis,” “probing,” “nasolacrimal obstruction,” and “infant lacrimal surgery.” Inclusion criteria covered infants aged 1–18 months presenting with persistent tearing, recurrent discharge, positive regurgitation test, or documented failure of conservative therapy. Data recorded included patient age, severity of symptoms, anatomical findings, type of obstruction, probing technique used, and postoperative response. Outcomes measured were symptom resolution time, recurrence frequency, need for secondary interventions, postoperative patency tests, and complication rates. Comparisons were made between early-probing groups and delayed-intervention groups to determine the impact of intervention timing.

Results:

Analysis of available data revealed that probing achieves high success rates, particularly in children treated before nine months of age. Early-probing cohorts exhibited rapid symptom disappearance, normalization of tear drainage, and minimal postoperative discomfort. Infants undergoing probing at later stages showed reduced success rates due to increasing duct rigidity and fibrotic tissue development. Cases with simple membranous obstruction demonstrated the highest success, whereas complex anatomical variants, canalicular stenosis, or bony anomalies exhibited relatively lower response rates and increased need for repeat procedures. Postoperative evaluation confirmed that the majority of children experienced complete resolution of mucopurulent discharge and elimination of recurrent infections within a short recovery period. Complication rates were low, with rare instances of minor bleeding or transient swelling. Comparative analysis showed significantly fewer secondary interventions in early-treated infants compared to older patients. Clinical observations indicate that high rates of resolution occur when probing is performed within the first year of life, with the most successful outcomes seen in infants younger than nine months. These patients exhibit rapid cessation of excessive tearing, disappearance of purulent reflux, and normalization of sac function. Obstructions of the simple membranous type respond most efficiently, whereas complex or anatomically irregular variants produce reduced success percentages and may require additional intervention. Delayed timing correlates with increasing tissue rigidity, more resistant membrane formation, and a greater need for secondary procedures such as intubation or balloon dilation. Post-procedure assessment reveals minimal adverse reactions, with only mild transient swelling noted in isolated cases. Overall, the majority of treated infants experience complete symptomatic recovery within a short period.

Discussion:

The positive outcomes associated with early probing validate its role as a standard therapeutic strategy for congenital dacryocystitis unresponsive to conservative measures. The high success rate in younger infants is attributed to greater ductal pliability, limited fibrosis, and improved membrane disruption efficiency. Delayed intervention, although still beneficial, demonstrates diminished effectiveness due to structural

rigidity and increased resistance at the distal obstruction site. Probing is minimally invasive, cost-effective, and associated with a rapid return to normal lacrimal function, making it a preferred technique in pediatric ophthalmology. Identifying appropriate candidates through clinical assessment, ensuring proper technique, and recognizing anatomical variations remain essential for optimizing outcomes. While most infants require only a single probing session, recurrence in complex cases highlights the need for alternative management strategies such as balloon dacryoplasty or silicone intubation.

The high effectiveness of probing underscores its importance as the primary surgical option for persistent congenital dacryocystitis. Its minimally invasive nature, combined with short operative duration and rapid improvement, makes it ideal for early management. Younger infants benefit most due to greater duct plasticity, which enhances membrane perforation and reduces resistance. Probing also decreases chronic inflammation risk, limits the development of fibrotic tissue, and minimizes long-term complications. Recognition of anatomical variations and selection of appropriate timing contribute significantly to procedural success. Although most cases require only one intervention, failure in complex obstruction highlights the necessity for alternative treatments. Nonetheless, its broad success supports widespread use in neonatal lacrimal pathology management.

Conclusion:

Probing of the lacrimal ducts in congenital dacryocystitis is a highly effective treatment, particularly when performed early, offering rapid symptom resolution, restoration of duct patency, and prevention of chronic complications. The technique’s reliability, low complication rate, and favorable anatomical responsiveness in younger patients establish it as the first-line surgical approach for persistent nasolacrimal duct obstruction. Incorporating timely evaluation, accurate diagnosis, and structured treatment planning ensures optimal results and reduces the necessity for additional procedures, contributing to improved pediatric ophthalmic care and long-term ocular health. Probing remains an effective and reliable treatment for congenital dacryocystitis unresponsive to conservative therapy, offering rapid symptom relief, restoration of normal tear drainage, and prevention of chronic disease progression. Early intervention provides optimal outcomes by utilizing the natural flexibility of the infant duct and reducing the likelihood of recurrent inflammation or structural alteration. Integrating timely probing into standard neonatal ophthalmic practice ensures improved functional recovery, reduces secondary surgical requirements, and supports long-term ocular health.

References:

1. Abduazizovich, Y. A., Abdurakhmanovich, B. S., Bakhodirovna, S. D., Batirovich, K. S., & Erkinovich, K. R. (2022). INTERRELATION OF FUNCTIONAL AND ANATOMICAL AND OPTICAL PARAMETERS OF THE EYE IN CONGENITAL MYOPIA. Web of Scientist: International Scientific Research Journal, 3(4), 582-590.
2. Abdurakhmanovich, B. S., Muratovna, K. A., Azizovich, Y. A., & Botirovich, K. S. Effectiveness Of Surgical Treatment Of High Myopia By Implantation Of Phakic Intraocular Lenses //European Journal of Molecular & Clinical Medicine, 7(03), 2020.
3. Бабаев, С. А., Кадирова, А. М., Юсупов, А. А., Бектурдиев, Ш. С., & Сабирова, Д. Б. Наш опыт хирургического исправления вторичного расходящегося косоглазия у детей // Точка зрения. Восток–Запад, (3), (2016). 124-126.
4. Бабаев, С. А., Кадирова, А. М., Садуллаев, А. Б., Бектурдиев, Ш. С., Салахиддинова, Ф. О., & Хамрокулов, С. Б. Эффективность операции факоэмульсификации с имплантацией интраокулярных линз при зрелых старческих катарактах // Вестник врача, (2017). (3), 23.

5. Бабаев, С. А., Кадирова, А. М., & Орипова, Е. Ч. ЭФФЕКТИВНОСТЬ ШОВНОГО МАТЕРИАЛА ПРЕМИЛЕНА В ХИРУРГИИ ВРОЖДЕННОГО БЛЕФАРОПТОЗА // ВЕСТНИК ВРАЧА, 20.
6. Бобоев, С. А., Кадирова, А. М., Исмоилов, Ж. Ж., Косимов, Р. Э., & Бобоев, С. С. ОПЫТ ТРАНССКЛЕРАЛЬНОЙ ЛАЗЕРНОЙ ФОТОКОАГУЛЯЦИИ ЦИЛИАРНОГО ТЕЛА У БОЛЬНЫХ С НЕОВАСКУЛЯРНОЙ ГЛАУКОМОЙ // In VOLGAMEDSCIENCE (2021). (pp. 430-432).
7. Долиев, М. Н., Тулакова, Г. Э., Кадырова, А. М., Юсупов, З. А., & Жалалова, Д. З. ЭФФЕКТИВНОСТЬ КОМБИНИРОВАННОГО ЛЕЧЕНИЯ ПАЦИЕНТОВ С ЦЕНТРАЛЬНОЙ СЕРОЗНОЙ ХОРИОРЕТИНОПАТИЕЙ // Вестник Башкирского государственного медицинского университета, (2016). (2), 64-66.
8. Жалалова, Д. З., Кадирова, А. М., & Хамракулов, С. Б. ИСХОДЫ ГЕРПЕТИЧЕСКИХ КЕРАТОУВЕИТОВ НА ФОНЕ ЛЕЧЕНИЯ ПРЕПАРАТОМ «ОФТАЛЬМОФЕРОН» В ЗАВИСИМОСТИ ОТ ИММУННОГО СТАТУСА ПАЦИЕНТОВ // МЕЖДИСЦИПЛИНАРНЫЙ ПОДХОД ПО ЗАБОЛЕВАНИЯМ ОРГАНОВ ГОЛОВЫ И ШЕИ, (2021). 103.
9. Жалалова, Д. З. Метод комбинированного лечения диабетической ретинопатии // Врач-аспирант, (2009). 37(10), 864-868.
10. Кадирова, А. М., Бобоев, С. А., & Хакимова, М. Ш. РАННЕЕ ВЫЯВЛЕНИЕ И ЛЕЧЕНИЕ СПАЗМА АККОМОДАЦИИ У ДЕТЕЙ // Форум молодых ученых, (2021) (5), 191-196.
11. Кадирова, А. М., Бобоев, С. А., & Хамракулов, С. Б. (2021). ЭФФЕКТИВНОСТЬ РЕТИНАЛАМИНА В ЛЕЧЕНИИ ВРОЖДЕННОЙ МИОПИИ. In VOLGAMEDSCIENCE (pp. 429-430).
12. Кадирова, А. М., Рузиев, Т. Х., & Хамракулов, С. Б. (2019). ОТДАЛЕННЫЕ РЕЗУЛЬТАТЫ АУТОПЛАСТИКИ КОНЪЮНКТИВАЛЬНЫМ ЛОСКУТОМ У БОЛЬНЫХ С КРЫЛОВИДНОЙ ПЛЕВОЙ. ТОМ–I, 235.
13. Кодирова А.М., Бабаев С.А., Каландаров Ф.У., Гаффаров Г.К. Эффективность дакриоцисториностомии биканальной интубацией слезной полости Силиконовыми трубчатыми путями // «На пути научных открытий». Материалы научно-практической конференции молодых ученых, 9 апреля, Ташкентское шоссе, 2013 г. стр. 231.
14. Кадирова А. М., Хамракулов С. Б., Хакимова М. Ш. ЛЕЧЕНИЕ СПАЗМА АККОМОДАЦИИ У ДЕТЕЙ //СОВРЕМЕННАЯ НАУКА: АКТУАЛЬНЫЕ ВОПРОСЫ И ПЕРСПЕКТИВЫ РАЗВИТИЯ. – 2021. – С. 231-236.
15. Мухамадиев, Р. О., Дехканов, Т. Д., Блинова, С. А., Юсупов, А. А., & Хамидова, Ф. М. Возрастные особенности кристаллизации слезы у здоровых лиц // ВЕСТНИК ВРАЧА, 26.
16. Мухамадиев, Р. О., Рахимова, Л. Д., Кадирова, А. М., & Хамидова, Ф. М. ХАР ХИЛ КЎЗ КАСАЛЛИКЛАРИДА КЎЗ ЁШЛАРИ КРИСТАЛЛОГРАФИЯСИ // МЕЖДИСЦИПЛИНАРНЫЙ ПОДХОД ПО ЗАБОЛЕВАНИЯМ ОРГАНОВ ГОЛОВЫ И ШЕИ, 123.
17. Сабирова, Д. Б., Юсупов, А. А., Искандаров, Ш. Х., Кадырова, А. М., & Тулакова, Г. Э. Клиническая оценка озонотерапии и криопексии у пациентов с герпетическим кератитом // Точка зрения. Восток–Запад, (2016). (1), 147-149.
18. Сабирова, Д. Б., Тулакова, Г. Э., & Эргашева, Д. С. Комплексное лечение диабетической макулопатии путем применения пептидного биорегулятора" Ретиналамин" и лазеркоагуляции сетчатки // Точка зрения. Восток-Запад, (2017). (2), 114-116.

19. Сабирова, Д. Б., Искандаров, Ш. Х., Косимов, Р. Э., Эргашева, Д. С., & Юсупов, А. А. Совершенствование лечения герпетических кератитов с использованием озона в виде газа через очки аппарата" Орион-си" // Российский общенациональный офтальмологический форум, (2015). 1, 159-163.

20. Сабирова, Д. Б., Облоёров, И. Х., & Хазратова, Д. Ф. КЛИНИКО-ЭПИДЕМИОЛОГИЧЕСКИЕ ОСОБЕННОСТИ ВЕСЕННЕГО КАТАРА И ЛЕЧЕНИЕ ИММУНОКОРРЕГИРУЮЩИМИ СРЕДСТВАМИ // НАУЧНЫЕ ИССЛЕДОВАНИЯ, (2019).52.

21. Саттарова, Х. С., Жалалова, Д. З., & Бектурдиев, Ш. С. Причины слепоты и слабовидения при сахарном диабете // Академический журнал Западной Сибири, (2011). (6), 27-28.

22. Тулакова, Г. Э., Сабирова, Д. Б., Хамракулов, С. Б., & Эргашева, Д. С. Отдалённые результаты ксеносклеропластики при миопии высокой степени // Научный форум. Сибирь, (2018). 4(1), 80-80.

23. Хамидова, Ф. М., Амриддинова, Ш. А., & Очилова, Н. Н. Ретиналамин в комплексном лечении больных с осложненной прогрессирующей миопией // Ответственный редактор, (2012). 3, 727.

24. Юсупов, А. А., Бобоев, С. А., Хамракулов, С. Б., Сабирова, Д. Б., & Косимов, Р. Э. Взаимосвязь функциональных и анатомо-оптических параметров глаза при врожденной близорукости // Вопросы науки и образования, (2020). (22 (106)), 44-53.

25. Юсупов, А., Хамракулов, С., Бобоев, С., Кадирова, А., Юсупова, Н., & Юсупова, М. (2021). АМЕТРОПИЯЛАРНИ ИНТРАОКУЛЯР ФАКИК ЛИНЗАЛАР БИЛАН КОРРЕКЦИЯЛАШ. Журнал стоматологии и краниофациальных исследований, 2(1),13-17.

26. Юсупов, А. А., Юсупова, Н. К., & Хамракулов, С. Б. Интроокулярная коррекция высокой анизометропии при косоглазии // Современные технологии в офтальмологии, (2020).(4), 251-252.

27. Юсупов, А. А., Кадирова, А. М., Бабаев, С. А., Очилова, Н. Н., Косимов, Р. Э., & Салахиддинова, Ф. О. Криопексия в комплексной терапии больных с неоваскулярной болящей глаукомой. // Российский общенациональный офтальмологический форум, (2015). 1, 196-198.

28. Юсупов, А. Ю., Салиев, М. С., Закирова, З. И., & Василенко, А. В. (1988). Некоторые особенности клинических проявлений глаукомы с относительно низким офтальмотонусом и различным уровнем артериального давления. Вестник офтальмологии, 104(5), 6-8.

29. Юсупов АЮ, Закирова ЗИ, Василенко АВ. Индекс интолерантности как прогностический ориентир при медикаментозном лечении глаукомы с низким внутриглазным давлением. InМеждунар. конф., посв (pp. 121-122).

30. Юсупов, А. А., Юсупова, М. А., Юсупова, Н. А., & Василенко, А. В. (2020). МИКРОИМПУЛЬС ТРАНСКЛЕРАЛЬНАЯ ЦИКЛОФОТОКОАГУЛЯЦИЯ (МТСЦФК) ПРИ ЛЕЧЕНИИ РЕФРАКТЕРНОЙ ГЛАУКОМЫ: ПРЕДВАРИТЕЛЬНЫЕ РЕЗУЛЬТАТЫ. In МИРОВАЯ ЭКОНОМИКА. ПРОБЛЕМЫ, ПЕРСПЕКТИВЫ, ИННОВАЦИИ (pp. 42-49).

31. Юсупов, А., Хамидова, Ф., & Василенко, А. (2022). Наш опыт сочетанного применения геля вирган и парабульбарных инъекций циклоферона при поверхностных формах герпетического кератита. Журнал вестник врача, 1(1), 215–216. извлечено от https://inlibrary.uz/index.php/doctors_herald/article/view/9103