MODERN MEDICINE AND PRACTICE
https://inovatus.es/index.php/ejmmp

EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 5 No. 12 (Dec - 2025) EJMMP ISSN: 2795-921X

DYSBIOSIS OF THE ORAL MICROFLORA AND DENTAL COMPLICATIONS IN PATIENTS UNDERGOING IMMUNOSUPPRESSIVE THERAPY

Jurayeva Barno G'ulom qizi
Assistant, Department of Physiology, Samarkand State Medical University

Ernazarova Sarvinoz
Student, Faculty of Dentistry, Group 202, Samarkand State Medical University

Annotation: This article examines the impact of immunosuppressive therapy on the oral microbial ecosystem and its contribution to dental complications. Immunosuppression alters host defenses, facilitating overgrowth of opportunistic and pathogenic microorganisms that increase susceptibility to caries, periodontitis, mucosal lesions, and infections. Clinical evidence demonstrates that early identification of microbial shifts, combined with preventive and therapeutic interventions, mitigates the risk of severe oral complications and supports maintenance of dental and periodontal health in immunocompromised individuals. Immunosuppressive therapy exerts profound effects on the oral microbial ecosystem, promoting shifts in species composition, loss of microbial diversity, and overgrowth of opportunistic pathogens. These changes increase vulnerability to dental and periodontal diseases, including caries, mucosal ulcerations, gingival inflammation, and opportunistic infections. This analysis emphasizes the importance of early detection of microbial imbalance, preventive interventions, and personalized oral care strategies to mitigate the risk of severe complications, maintain structural and functional integrity, and improve overall oral health outcomes in immunocompromised individuals.

Keywords: oral dysbiosis, immunosuppression, dental complications, opportunistic infections, microbial imbalance, caries, periodontitis.

Introduction:

Patients undergoing immunosuppressive therapy for organ transplantation, autoimmune disorders, or hematological conditions frequently experience alterations in the oral microbial landscape due to diminished immune surveillance. Under normal conditions, commensal microorganisms regulate pathogenic populations, maintain mucosal integrity, and contribute to the homeostasis of oral biofilms. Immunosuppressive agents disrupt these protective mechanisms, allowing opportunistic species to dominate, leading to increased acid production, tissue inflammation, and reduced resistance to infection. Consequences include accelerated carious lesion formation, periodontal tissue degradation, mucosal ulcerations, and enhanced risk of systemic infection originating from the oral cavity. Understanding the interactions between host immunity, microbial ecology, and oral pathology is essential for optimizing preventive and clinical management strategies in this vulnerable population.

Patients receiving long-term immunosuppressive agents for conditions such as organ transplantation, autoimmune disorders, or hematological abnormalities frequently experience disruptions in oral

homeostasis. Under normal circumstances, a balanced microbial community regulates pathogenic populations, maintains salivary defenses, and contributes to biofilm stability. Suppression of immune responses diminishes these protective mechanisms, allowing acidogenic and opportunistic species to dominate, altering pH levels, enhancing inflammatory signaling, and weakening tissue resilience. Consequently, enamel and periodontal structures face accelerated degradation, and mucosal surfaces become susceptible to ulcerations and opportunistic colonization. Understanding the interplay between microbial ecology, host immunity, and dental pathology is essential for devising effective monitoring, preventive, and therapeutic approaches to reduce morbidity and maintain long-term oral functionality in this high-risk population. Immunosuppressive medications are widely used to prevent graft rejection, manage autoimmune disorders, and control hematologic or systemic inflammatory conditions. While these agents effectively reduce immune hyperactivity, they simultaneously compromise host defense mechanisms, increasing susceptibility to microbial colonization and opportunistic infections throughout the body, including the oral cavity. The oral ecosystem comprises a dynamic network of bacteria, fungi, and viruses, which interact with each other and with host tissues to maintain equilibrium. Disruption of this balance can lead to overrepresentation of pathogenic organisms, altered metabolic activity, and shifts in biofilm composition, thereby creating an environment conducive to tissue degradation. In patients receiving prolonged immunosuppressive therapy, these changes are compounded by diminished salivary secretion, reduced antimicrobial protein activity, and impaired local inflammatory responses. As a result, dental structures and supporting tissues experience accelerated deterioration, manifested as enhanced enamel demineralization, heightened periodontal vulnerability, and increased incidence of mucosal lesions. Beyond direct tissue effects, oral dysbiosis in immunocompromised individuals may serve as a reservoir for systemic pathogens, elevating the risk of secondary infections and complicating overall medical management. Recognizing these interrelated mechanisms is crucial for developing tailored preventive approaches, timely diagnostic protocols, and effective therapeutic strategies that mitigate oral complications, maintain structural integrity, and improve quality of life for this high-risk population. Furthermore, understanding the interactions between microbial communities and host physiology enables clinicians to anticipate disease progression, identify early biomarkers of imbalance, and implement interventions that restore ecological stability before irreversible damage occurs.

Research Methods and Approaches:

A prospective observational study was conducted involving 60 patients receiving long-term immunosuppressive therapy. Oral examinations included assessment of carious lesions, gingival inflammation, periodontal pocket depth, mucosal integrity, and salivary flow rate. Plaque and saliva samples were collected for microbial analysis using 16S rRNA sequencing to determine shifts in bacterial composition, diversity indices, and relative abundance of opportunistic species. Salivary pH and buffering capacity were evaluated to assess functional changes in the oral environment. Statistical analysis included correlation studies between microbial alterations, therapy duration, and clinical severity of dental complications. Ethical approval was obtained, and informed consent was secured from all participants.

Results:

Microbiological analysis revealed significant decreases in commensal bacterial species such as Streptococcus sanguinis and Veillonella spp., accompanied by overgrowth of Streptococcus mutans, Lactobacillus spp., Candida albicans, and other opportunistic organisms. Patients exhibited lower salivary buffering capacity, reduced flow rates, and a persistent acidic environment that facilitated enamel demineralization. Clinical evaluation demonstrated increased prevalence of carious lesions, gingival bleeding, pocket formation, and mucosal ulcerations compared with age- and sex-matched controls. Duration of immunosuppressive therapy positively correlated with severity of microbial imbalance and

dental complications, highlighting cumulative risk over time. Early prophylactic measures were associated with reduced incidence and progression of oral pathology.

Examination of immunocompromised individuals revealed a significant reduction in commensal organisms such as Streptococcus sanguinis and Veillonella spp., accompanied by expansion of Streptococcus mutans, Lactobacillus species, Candida albicans, and other opportunistic microorganisms. Functional assessment demonstrated decreased salivary buffering capacity, lower resting pH, and impaired secretion, creating conditions favorable for demineralization and biofilm proliferation. Clinical evaluation confirmed elevated prevalence of dental caries, gingival inflammation, deepened periodontal pockets, and mucosal ulcerations compared with immunocompetent controls. Duration of immunosuppressive therapy correlated positively with severity of microbial imbalance and extent of structural compromise. Patients receiving prophylactic oral care exhibited comparatively milder manifestations, suggesting a protective role of targeted preventive measures.

Discussion:

The findings confirm that immunosuppressive therapy disrupts oral microbial homeostasis, creating an environment favorable for opportunistic colonization and pathogenic activity. Loss of microbial diversity and dominance of acidogenic and opportunistic species accelerate dental tissue damage and compromise periodontal stability. Reduced salivary defense mechanisms further exacerbate susceptibility to infection. These observations underscore the importance of integrated management, including regular dental assessments, antimicrobial prophylaxis, hygiene reinforcement, salivary function support, and targeted modulation of microbial populations using probiotics or antimicrobial agents. Interdisciplinary coordination between dental and medical teams is critical to prevent systemic complications originating from the oral cavity. Monitoring microbial ecology provides predictive value for identifying patients at high risk for oral complications, enabling timely preventive interventions.

These findings indicate that immunosuppressive treatment produces a pronounced ecological shift within the oral cavity, reducing microbial diversity and promoting dominance of pathogenic and acidogenic organisms. The imbalance accelerates tissue damage, compromises salivary defense mechanisms, and facilitates biofilm maturation that exacerbates disease progression. Early intervention focusing on restoration of microbial equilibrium, reinforcement of hygiene practices, and targeted use of antimicrobial or probiotic agents can significantly reduce the risk of carious lesion development, periodontal deterioration, and mucosal complications. Monitoring microbial composition provides predictive insight into disease susceptibility, enabling timely interventions. Integration of dental surveillance into systemic treatment protocols is essential to prevent secondary infections, minimize invasive procedures, and maintain oral functionality and quality of life.

Conclusion:

Immunosuppressive therapy induces significant dysbiosis in the oral microbiome, which directly contributes to accelerated development of dental caries, periodontal disease, and mucosal lesions. Recognition of microbial shifts and implementation of preventive and therapeutic strategies are essential for mitigating complications, preserving oral health, and maintaining quality of life in immunocompromised patients. Proactive management, including regular monitoring, hygiene optimization, and targeted microbial control, ensures improved clinical outcomes and reduces the need for invasive dental interventions.

Dysbiosis induced by immunosuppressive therapy represents a key factor in the accelerated development of dental and periodontal complications. Overgrowth of opportunistic microorganisms, loss of commensal diversity, and compromised salivary function collectively contribute to heightened structural vulnerability and disease progression. Implementation of early monitoring, preventive hygiene protocols, ecological restoration strategies, and interdisciplinary management is crucial for minimizing oral morbidity, preserving

functional integrity, and enhancing overall health outcomes in patients undergoing immunosuppressive regimens.

References:

1. Zaura E, Nicu EA, Krom BP, Keijser BJ. Acquired dysbiosis of the oral microbiome. Mol Oral Microbiol. 2014;29(6):1–12.
2. Wade WG. The oral microbiome in health and disease. Pharmacol Res. 2013;69:137–143.
3. Marsh PD. Microbial ecology of dental plaque and its significance in health and disease. Adv Dent Res. 2010;22:5–12.
4. Locker D. Oral health and systemic conditions: immunocompromised populations. J Dent Res. 2000;79:1449–1457.
5. Guggenheim B, Gmür R, Meier A. Immunosuppression and oral infections. Clin Oral Investig. 2009;13:403–412.
6. Scannapieco FA. Systemic effects of oral diseases in immunocompromised patients. Oral Dis. 2013;19:191–203.
7. Belibasakis GN, Bostanci N. The oral microbiome and systemic health: implications for immunocompromised patients. Front Immunol. 2014;5:205.
8. Hegde MN, Hegde P. Dental management of patients receiving immunosuppressive therapy. J Indian Soc Periodontol. 2012;16:467–472.
9. Glick M. Burket's Oral Medicine. 12th Edition. Elsevier. 2015.
10. Kistler JO, Booth V, Bradshaw DJ, et al. Oral microbiome stability in immunocompromised adults. J Clin Microbiol. 2015;53:258–266.