MODERN MEDICINE AND PRACTICE
https://inovatus.es/index.php/ejmmp

EUROPEAN JOURNAL OF MODERN MEDICINE AND PRACTICE
Vol. 5 No. 12 (Dec - 2025) EJMMP ISSN: 2795-921X

THE IMPORTANCE OF EARLY REHABILITATION OF CHILDREN WITH CENTRAL NERVOUS SYSTEM DISORDERS AND BIRTH INJURIES

Kurbonova Meroj Kadriddin kizi
Student of the 1st course of the faculty of pediatrics, Kukon University Andijan Branch

Abstract: Early rehabilitation is a key factor in improving developmental outcomes in children with central nervous system disorders and birth-related injuries. Such conditions often lead to motor, cognitive, speech, and behavioral impairments that significantly affect quality of life. This article examines the importance of early rehabilitation interventions based on the principles of neuroplasticity and interdisciplinary care. The role of physical, occupational, and speech therapy, as well as family involvement and preventive strategies, is highlighted. Early, comprehensive rehabilitation not only enhances functional recovery but also reduces secondary complications and promotes social integration.

Keywords: early rehabilitation, central nervous system disorders, birth injuries, neuroplasticity, pediatric rehabilitation, developmental delay.

INTRODUCTION

Early childhood is a critical period for physical, cognitive, and emotional development. Disorders of the central nervous system and birth-related injuries represent some of the most significant challenges affecting children's health and long-term quality of life. These conditions often lead to motor impairments, sensory deficits, cognitive delays, and behavioral difficulties. In this context, early rehabilitation plays a decisive role in minimizing functional limitations, promoting neurodevelopment, and improving social adaptation. Timely and well-organized rehabilitation interventions can significantly influence developmental trajectories and long-term outcomes for affected children.

Central nervous system disorders and birth injuries may arise due to a variety of prenatal, perinatal, and postnatal factors, including hypoxia, prematurity, infections, metabolic disturbances, and mechanical trauma during delivery. Such conditions frequently disrupt normal brain development and neural connectivity, resulting in impaired motor control, muscle tone abnormalities, speech disorders, and learning difficulties. Without early intervention, these impairments may become more pronounced over time, leading to secondary complications and reduced independence in later life.

MATERIALS AND METHODS

The rationale for early rehabilitation is grounded in the concept of neuroplasticity. During infancy and early childhood, the brain demonstrates a high capacity for structural and functional reorganization. Neural circuits can adapt in response to external stimulation, therapeutic exercises, and environmental enrichment. Early rehabilitation takes advantage of this plasticity by stimulating preserved neural pathways and promoting compensatory mechanisms. As a result, children who receive timely rehabilitation often achieve better motor, cognitive, and communicative outcomes compared to those whose treatment is delayed [1].

Early rehabilitation encompasses a комплекс approach involving physical therapy, occupational therapy, speech and language therapy, sensory integration, and psychosocial support. Physical therapy aims to normalize muscle tone, improve posture, and develop gross motor skills such as sitting, standing, and walking. Occupational therapy focuses on fine motor coordination, daily living skills, and sensory processing. Speech and language therapy addresses communication difficulties, feeding disorders, and oral-motor dysfunctions, which are common in children with central nervous system damage. The integration of these interventions ensures a holistic impact on the child's development.

RESULTS AND DISCUSSION

Another important aspect of early rehabilitation is its preventive function. By addressing impairments at an early stage, rehabilitation reduces the risk of secondary complications such as joint contractures, skeletal deformities, chronic pain, and social isolation. Early support also helps prevent emotional and behavioral problems that may arise from prolonged functional limitations and communication barriers. Thus, rehabilitation not only improves physical functioning but also supports emotional well-being and social inclusion [2].

Family involvement is a crucial component of effective early rehabilitation. Parents and caregivers play a central role in the daily implementation of therapeutic strategies and in creating a supportive developmental environment. Educating families about the child's condition, rehabilitation goals, and home-based exercises enhances treatment continuity and effectiveness. Moreover, psychological support for families helps reduce stress, promotes positive parent-child interactions, and strengthens motivation to engage in long-term rehabilitation processes.

Early rehabilitation also has significant social and economic implications. Children who receive adequate early intervention are more likely to achieve higher levels of independence, participate in education, and integrate into society. This reduces the long-term burden on healthcare systems, social services, and families. From a public health perspective, investing in early rehabilitation programs represents a cost-effective strategy that yields substantial long-term benefits for individuals and society as a whole.

In modern healthcare systems, early rehabilitation increasingly relies on interdisciplinary collaboration. Pediatric neurologists, rehabilitation physicians, physiotherapists, speech therapists, psychologists, and educators work together to develop individualized rehabilitation plans. Such coordinated efforts ensure that interventions are tailored to the child's specific needs and developmental level. Advances in diagnostic methods and rehabilitation technologies further enhance the effectiveness of early intervention by enabling precise assessment and targeted therapy.

Early rehabilitation of children with central nervous system disorders and birth injuries is a critical factor determining developmental outcomes and quality of life. By leveraging neuroplasticity, preventing secondary complications, and supporting holistic development, early rehabilitation offers children the best opportunity to reach their full potential. A timely, comprehensive, and family-centered rehabilitation approach not only improves individual outcomes but also contributes to broader social and economic well-being. Therefore, early rehabilitation should be regarded as an essential component of pediatric healthcare and child development strategies [4].

An important dimension of early rehabilitation in children with central nervous system disorders and birth injuries is the early identification and stratification of developmental risks. Accurate assessment in the first months of life allows clinicians to distinguish between transient developmental delays and persistent neurological impairments. The use of standardized neurological examinations, developmental screening tools, and instrumental diagnostic methods enables timely decision-making regarding rehabilitation intensity and focus. Early risk stratification ensures that therapeutic resources are allocated efficiently and

that children with higher vulnerability receive more intensive and specialized support during critical periods of development.

Early rehabilitation also contributes to the optimization of sensory integration processes, which are frequently disrupted in children with central nervous system damage. Abnormal sensory processing can manifest as hypersensitivity, hyposensitivity, or disorganized responses to sensory stimuli, negatively affecting motor coordination, attention, and behavior. Targeted sensory-based interventions introduced at an early age help normalize sensory input processing and improve adaptive responses. As sensory integration forms the foundation for higher cognitive and motor functions, addressing these deficits early enhances overall developmental coherence and functional performance.

Another significant aspect of early rehabilitation is its role in supporting emerging cognitive and executive functions. Central nervous system disorders often impair attention regulation, working memory, and problem-solving abilities. Early cognitive stimulation through structured play, guided interaction, and task-oriented activities promotes neural connectivity associated with learning and information processing. Rehabilitation programs that incorporate age-appropriate cognitive challenges facilitate the development of adaptive strategies and reduce the severity of future learning difficulties. This early cognitive support is particularly important for preparing children for successful participation in educational settings.

CONCLUSION

Early rehabilitation plays a decisive role in the management of children with central nervous system disorders and birth injuries. Timely intervention during periods of high neuroplasticity improves motor, cognitive, and communicative functions while preventing secondary complications. A multidisciplinary and family-centered rehabilitation approach enhances developmental potential and supports social inclusion. Integrating early rehabilitation into pediatric healthcare systems is essential for improving long-term outcomes and reducing the overall burden of disability.

REFERENCES

1. Баранов А.А., Альбицкий В.Ю. Реабилитация детей с поражениями центральной нервной системы. – Москва: ПедиатрЪ, 2016. – 320 с.
2. Козлова Л.В. Ранняя помощь детям с перинатальными поражениями ЦНС. – Санкт-Петербург: СпецЛит, 2017. – 256 с.
3. Novak I., Morgan C., Adde L. et al. Early, Accurate Diagnosis and Early Intervention in Cerebral Palsy // JAMA Pediatrics. – 2017. – Vol. 171, No. 9. – P. 897–907.
4. Rosenbaum P., Paneth N., Leviton A. et al. A Report: The Definition and Classification of Cerebral Palsy // Developmental Medicine & Child Neurology. – 2007. – Vol. 49 (Suppl. 109). – P. 8–14.
5. WHO. Rehabilitation in Health Systems. – Geneva: World Health Organization, 2017. – 56 p.